

Molecular and Serological Studies of Iris Yellow Spot Virus on Onion Plants in Iran

¹Neda Rafizadeh, ¹Behrouz Jafarpour, ²Mehdi Nasr Isfahani and ¹Mahrokh Falahati Rastegar

¹Department of Plant Protection, Ferdowsi University of Mashhad, Iran

²Department of Plant Protection, Isfahan Research Center for Agricultural Science,
Natural Resources Isfahan, Iran

Abstract: The Iris Yellow Spot Virus (IYSV) is one of the diseases of Iris plants in Iran, occurring annually and causing considerable losses. In this manuscript, the molecular and serological studies of the disease were taken placed for the 1st time in Iran. For this purpose, 435 samples with symptoms of chlorotic, necrotic and diamond shape lesion were collected from onion fields and 142 from ornamental plants. The infected plants were transferred into the laboratory for DAS-ELISA tests and the sap of positive plants inoculated to 4 indicator host plants *Nicotiana rustica* (with leaf deforming and systemic chlorotic and necrosis), *N. benthamiana*, *N. clevelandii*, *N. tabacum* var. *samson* (with systemic chlorotic and necrosis) in greenhouse conditions. The inoculated indicator plants tested for DAS-ELISA then sap of positive indicator plants inoculated to 4 onion genotypes, neishabour yellow, neishabour white, dargaz red and isfahan red (Dorche). Also for genetical analysis, the RNAs were extracted by PEG6000 precipitation and RNXTM (plus) kit. In RT-PCR tests, specific primers designed for nucleoprotein gene amplified 181 and 139 bp fragments. It was found that all of the onion fields were infected with the virus in various degrees, out of which IYSV was detected in 107 samples of onion, 7 samples of chrysanthemum flowers and a sample of Iris flowers.

Key words: Iris yellow spot virus, DAS-ELISA, onion, RT-PCR, Iris flowers, Iran

INTRODUCTION

Iris yellow spot virus was first reported affecting onion inflorescence in Southern Brazil in 1981. Then after the disease was not seen again until 1994. When it was detected in North-Eastern Brazil, the disease was characterized by symptoms of chlorotic and necrotic eye-like or diamond shaped lesions inflorescence (Gent *et al.*, 2006). Cortez *et al.* (1998) described IYSV in the Netherlands as a new Tospovirus naturally infecting Iris hollandica in the field and leek in the greenhouse. Near the same time, Gera *et al.* (1998) reported that IYSV on onion in Israel. Kritzman *et al.* (2000) reported natural IYSV infection of lisianthus (*Eustoma russellianum*) grown in the field in Israel.

IYSV is now known to occur on onion in the following locations: 1999, India; 2000, Slovenia; 2002, Colorado (USA); 2003, Australia and Italy; 2004, Japan, Georgia (USA), New Mexico (USA) and Washington (USA); 2005, Chile, Peru, Spain, Tunisia, central Oregon (USA) and 2006, Reunion Island, Guatemala, Texas (USA) and New York (USA) (Gent *et al.*, 2006). In 2007, Mumford *et al.* (2008) reported this virus on *E. grandiflorum*. Gawande *et al.* (2010) reported this virus on garlic in India for the 1st time. Ghotbi *et al.* (2005)

reported this virus on *Plargonium hortorum*, *Rosa* sp., *Cycas* sp. and *Scindapsus* sp. IYSV is one of members of family Bunyaviridae genus Tospovirus. The viruses of this genus have enveloped isometric particles (80-110 nm) which contain virus-encoded Glycoproteins (G1/G2). The genome of Tospoviruses consists of three ssRNA segments denoted L RNA, M RNA and S RNA. The M RNA encodes the Glycoproteins G1/G2 (Silva *et al.*, 2001). Onion thrips, Thrips tabaci (Order, Thysanoptera, Family, Thripidae) is a worldwide pest of vegetable crops. It ranges from tropical and subtropical areas into the temperate regions, *T. tabaci* is an important pest of field and greenhouse crops all around the world. It causes damages directly by feeding and indirectly by transmitting Tomato Spotted Wilt Virus (TSWV) but only the larval stage can transmit this virus (Pourian *et al.*, 2009). IYSV is transmitted by thrips in a persistent manner. Once a thrips has acquired the virus, it can transmit the virus for the remainder of its lifetime (Crowe and Pappu, 2005). Viruliferous thrips emigrating from harvested onion fields into nonharvested ones may be increasing the primary spread of IYSV in late-harvested onions. Managing *T. tabaci* populations before harvest and manipulating the spatial arrangement of fields based on harvest date could mitigate the spread of IYSV (Hsu *et al.*,

2010). The disease was characterized by symptoms of chlorotic and necrotic eye-like or diamond shaped lesions on scapes (Pozzer *et al.*, 1994). Symptoms appeared as chlorotic or necrotic, straw-colored to white, dry, elongate or spindle-shaped lesions along the scape with lesions frequently more numerous at mid to lower portions of the scape. In some lesions, an island of green tissue developed in the center of the necrotic area. When lesions became large and numerous, they coalesced often completely girdling the scape. This weakened the scape, causing the seed head to collapse and topple over. Patterns of disease incidence in fields or locations were not apparent, nor was there any association with host genotype or cultural practices. Estimated yield losses in individual fields ranged from insignificant to nearly 60% (Mohan and Wilson, 1989). Reports of the economic losses caused by IYSV in onion range were from high in Israel and Brazil (Smith *et al.*, 2006) to minimal in the Netherlands (Cortez *et al.*, 1998). Symptoms on leaves that could be confused with those caused by fungal or bacterial diseases. Additionally, the necrotic areas resulting from IYSV infection could be colonized by secondary invaders such as *Stemphylium* sp. or *Alternaria* sp., leading to inaccurate diagnosis. (Pappu *et al.*, 2008). The disease like as Cladosporium leaf spot (Evans *et al.*, 2009). Most important of indicator plants for this virus are *Nicotiana benthamiana* and *N. rustica* with systemic symptoms (Chatzivassiliou *et al.*, 2000). IYSV does not appear to be seed-borne or seed-transmitted in onion (Bulajic *et al.*, 2009).

MATERIALS AND METHODS

Sampling: An intensive survey was conducted during 2008-2009 to determine the molecular and serological studies of the IYSV on ornamentals and onion plants. In 2008, during an initial survey, 435 samples of onion and 142 samples of ornamental plants were collected. Samples of ornamentals comprised of leaves from different parts of each plant exhibited various symptom types suggestive of Tospovirus infection, such as yellow or necrotic spots on leaves and tip necrosis. Symptomless onion leaf and scape samples were collected together with those showing symptoms of elongated, chlorotic, necrotic, oval chlorotic, diamond shape lesion or necrotic lesions. Both fresh and frozen (-80°C) samples were tested for IYSV presence by Enzyme Linked Immunosorbent Assay (ELISA).

Serological testing: Standard DAS-ELISA was performed. Samples were tested for the presence of IYSV utilizing a Double Antibody Sandwich (DAS)-ELISA Method of (Clark and Adams, 1977) serological reagents

against IYSV used was from the DSMZ (Braunschweig, Germany). Absorbance at 405 nm was measured with an ELISA microplate reader. Samples were considered positive if the absorbance value was ≥ 2 times the absorbance of the negative control.

Mechanical inoculation: *Nicotiana benthamiana*, *N. rustica*, *N. tabacum* var. *samson* and *N. clevelandii* were mechanically inoculated with infected onion sap, using 0.01 M potassium phosphate buffer, pH 7.0, containing 0.2% sodium sulfite and 0.01M 2-mercaptoethanol (Mandal *et al.*, 2001). A total of 4 plants of each experimental species were inoculated and the bioassay was repeated three times. After inoculation, the plants were sprayed with distilled water and kept in an insect-proof greenhouse at a temperature of 22-25°C and were inspected regularly for symptom development. About 3 weeks after inoculation both symptomatic and asymptomatic plants were assayed by DAS-ELISA to confirm IYSV presence and to detect symptomless infections. Some of the ELISA positive leaves were further tested using Reverse Transcription-Polymerase Chain Reaction (RT-PCR). Also in order to investigate virus transmission to onion plants, 4 onion genotypes (yellow of Neishabour, red of Dorche Isfahan, red of Dargaz and white of Neishabour) were mechanically inoculated with infected nicotine sap using 0.01 M potassium phosphate buffer, pH 7.0, containing 0.2% sodium sulfite and 0.01 M 2-mercaptoethanol (Mandal *et al.*, 2001). Total of 4 onion plants of each experimental genotypes were inoculated and the bioassay was repeated 3 times. Plants were kept in an insect-proof greenhouse at a temperature of 22-25°C and were inspected regularly for symptom development. Three weeks after inoculation, both symptomatic and asymptomatic plants were assayed by DAS-ELISA to confirm IYSV presence and to detect symptomless infections.

RNA extraction: Viral RNA extraction from systemically infected nicotine and onion plants were performed. Two methods were used:

- Sedimentation with PEG 6000 base on Smith *et al.* (2006) was done
- Extraction with plus (RNX™) mixture used from Cinagen company was done

Electrophoresis of extracted RNA in 1% agarose gel: In order to investigate of RNA quality 3 μ L extracted RNA with 1 μ L of color buffer were subjected to electrophoresis in a 1% agarose gel, stained with 0.5 μ g mL⁻¹ ethidium bromide and photographed under UV illumination.

Table 1: Primers used for Iris yellow spot virus detection

Primers	Direction	Sequence of primers (5'-3')	Location	(bp)
IYSV-F1	Forward	GAGATGTGGATGTGGTGATTG	Coat protein	139
IYSV-R1	Reverse	GTCTTGTAATGCCTGCTCTGT		
IYSV-F2	Forward	TAGGGTGAAACCGTCAGAAA	Coat protein	181
IYSV-R1	Reverse	TGTCTTGTAATGCCTGCTC		

Table 2: RT-PCR mixture

Quantity	Material
2.5 µL	10×Buffer
0.5 µL	Dntp
1 µL	MgCl ₂
0.3 µL	Taq DNA polymerase
1 µL	Reverse primer
5 µL	cDNA
13.7 µL	Sterile water

Table 3: PCR program

Cycle no.	Time	Temperature (°C)
Stage 2-36 cycle	3 min	94
	30 sec	94
	30 sec	61
	20 sec	72
	30 sec	72
	10 min	4

Primers: The primers used for the cDNA synthesis were either designed based on S RNA of coat protein gene of IYSV previously reported by Ward *et al.* (2008). Primers were synthesis by Bioneer Company (Table 1).

Amplification of cDNA with Polymerase Chain Reaction (PCR):

First strand cDNA was primed on total plant RNA using specific reverse primers using Accupower TMRT Premix kit (Bioneer Company).

RT-PCR: RT-PCR was carried out with the Accupower TMRT Premix kit (Bioneer Company) according to the manufacturer's instructions. The RT-PCR reaction mixture included mixture included 2.5 µL 10×Buffer, 0.5 µL dntp, 1 µL MgCl₂, 0.3 µL Taq DNA Polymerase, 1 µL Forward Primer, 1 µL Reverse Primer, 5 µL cDNA in a final volume of 25 µL (Table 2). Cycling conditions were as follows (Table 3).

Agarose gel electrophoresis: Amplified products were analyzed by 1.7% agarose gel electrophoresis, stained with ethidium bromide and visualized under UV transilluminator.

RESULTS

Virus symptoms: Onion plants with chlorotic or necrotic and diamond-shaped lesions on leaves and especially on scapes (Fig. 1 and 2) were observed in onion fields. Affected plants were spread across the field with very high disease incidence.



Fig. 1: Symptoms of Iris yellow spot virus in onion field diamond shape lesion (on leaves)



Fig. 2: Symptoms of Iris yellow spot virus in onion field diamond shape lesion (on scapes)

Disease occurrence was associated with a high population of *T. tabaci*. Also Iris and Chrysanthemum flowers with necrotic lesions on leaves and especially on scapes were observed in greenhouse.

Virus detection and result of ELISA test: Of the 577 samples assayed, 115 reacted positively in DAS-ELISA. 435 samples with symptoms of chlorotic, necrotic and diamond shape lesion were collected from onion fields and 142 from ornamental plants (Rose, Gladiol, Iris, Plargonium, Chrysanthemum, Begonia, Petonia and Carnation).

Table 4: Infected plants and percent of infected plants

Plants	Total no.	Infected plants	Infected plants (%)
Onion	435	107	24.59
Iris	8	1	---
Ornamental plants	142	-	5.63
Chrysanthemum	7	7	---




Fig. 3: Percent of infected plants in onion and ornamental plants

IYSV was detected by ELISA in 107 samples of onion (24.59%), 7 samples of chrysanthemum flowers and a sample of Iris flowers (Table 4 and Fig. 3).

Mechanical incubation

Mechanical incubation of indicator plants: Different plant species from a botanical family (Nicotiana) were inoculated from selected ELISA-positive plants with Iris yellow spot virus. After inoculation, the plants were sprayed with distilled water and kept in an insect-proof greenhouse at a temperature of 22-25°C and were inspected regularly for symptom development. On inoculated leaves of plants symptoms developed within 2 weeks post inoculation. Symptom descriptions on indicator test plants for these viruses are given:

- *N. benthamiana* and *N. clevelandii* showed chlorosis and systemic necrosis
- *N. tabacum* var. *samson* gave chlorosis and systemic necrosis symptoms
- *N. rustica* gave leaf deformation and systemic chlorosis and necrosis

About 3 weeks after inoculation, both symptomatic and asymptomatic plants were assayed by DAS-ELISA to confirm IYSV presence and to detect symptomless infections (Table 5 and Fig. 4, 5).

Mechanical incubation of onion genotype: Four onion genotypes (Yellow of Neishabour, Red of Dorche Isfahan,

Table 5: Percent of infected indicator plants

Plant cultivar	No. of plants	Infected plants	Infection (%)
<i>Nicotiana tabacum</i>	15	7	46.6
var. <i>samson</i>			
<i>N. rustica</i>	10	4	40.0
<i>Nicotiana benthamiana</i>	13	6	46.1
<i>Nicotiana clevelandii</i>	16	10	62.5

Fig. 4: Symptoms of IYSV in *N. rustica* deforming leaves, systemic chlorosis and necrosisFig. 5: Symptoms of IYSV in *N. clevelandii*: Chlorosis and systemic necrosis

Red of Dar gaz and White of Neishabour) were mechanically inoculated with infected nicotine sap. Onion plants were inoculated and were kept in an insect-proof

Table 6: Results of DAS-ELISA test of cultivars of onion

Name of cultivar	No. of plants	Infected plants	Infection (%)
Neishabour yellow	15	7	46.6
Isfahan red (Dorche)	12	6	50.0
Dargaz red	13	3	23.0
Neishabour white	17	1	5.8

Fig. 6: Symptoms of IYSV in *N. tabaccum* var. *samson*: Chlorosis and sustemic necrosisFig. 7: Symptoms of IYSV in *N. Benthamiana*: Chlorosis and systemic necrosis

greenhouse at a temperature of 22-25°C and were inspected regularly for symptom development. On inoculated leaves of onion plants symptoms (chlorotic and necrotic lesions on leaves and especially on scapes) developed within 2 weeks post inoculation. About 3 weeks after inoculation both symptomatic and asymptomatic plants were assayed by DAS-ELISA to confirm IYSV presence and to detect symptomless infections (Table 6 and Fig. 6 and 7).

RNA electrophoresis: Viral RNA extraction from systemically infected nicotine and onion plants were performed. In order to investigate of RNA quality 3 µL extracted RNA with 1 µL of color buffer were subjected to





Fig. 8: Analysis of RNA of iris yellow spot virus on 1% agarose gel

Fig. 9: Results of RT-PCR for sample amplified with primers IYSV-F1/IYSV-R1 number is *I. ladear*

electrophoresis in a 1% agarose gel, stained with 0.5 µg mL⁻¹ ethidium bromide and photographed under UV illumination (Fig. 8).

Result of RT-PCR test and diagnose of virus: First strand cDNA was primed on total plant RNA using specific reverse primers using Accupower TMRT Premix kit (Bioneer Company). RT-PCR was carried out with the Accupower TMRT Premix kit (Bioneer Company) according to the manufacturer's instructions. Amplified products were analyzed by 1.7% agarose gel electrophoresis, stained with ethidium bromide and visualized under a UV transilluminator (Fig. 9). For having high quality Touchdown PCR Method used for delete the additional sections in primer 181 bp (Fig. 10). The primers were able to amplify the amplicon of expected size. RT-PCR with primers IYSV-F1/IYSV-R1, amplified fragments of the expected size (139 bp) from samples infected with IYSV (Fig. 9). Similar results were obtained with primers IYSV-F2/IYSV-R2 that amplified a fragment of 181 bp (Fig. 10).




Fig. 10: Results of RT-PCR for sample amplified with primers IYSV-FI/IYSV-RI number is *I. ladear*

DISCUSSION

In this study, the molecular and serological studies of the disease were taken placed. Symptoms caused by IYSV in inoculated onion plants in greenhouse (chlorotic and necrotic lesions on leaves and especially on scapes) were very similar to those reported by Mohan and Wilson (1989). This study confirmed that mechanical transmission of IYSV from infected onion samples naturally is possible. Sometime this virus caused symptomless infection of *N. benthamiana*, *N. clevelandii*, *N. tabacum* var. *samson* and onion plants while IYSV always caused symptomatic infection (local lesions) of *N. tabacum*. Symptoms caused by IYSV in *N. rustica* (leaf deformation and systemic chlorosis and necrosis) were very similar to those reported by Pozzer *et al.* (1999). Symptoms caused by IYSV in *N. clevelandii* (systemic chlorose) were similar to those reported by Ravi *et al.* (2006). *N. benthamiana* and *N. clevelandii* showed chlorosis and systemic necrosis. *N. clevelandii* had maximum infection so suggest that use this cultivar in experience methods of this virus in spite of Chatzivassiliou *et al.* (2000) reported that *N. benthamiana* and *N. rustica* was most important indicator for this virus. Symptom of indicator plants that inoculated with virus and results of DAS-ELISA test and RT-PCR test indicate that infection of IYSV is becoming an increasingly important constraint to onion production. Rapid and

accurate diagnostic tools and methods are essential to better understand the epidemiology of the virus and to devise effective management strategies. Standard RT-PCR was more sensitive and rapid than DAS-ELISA. Comparison of four cultivar of onion virus transmission in greenhouse showed that a maximum rate of transmission achieved in red of Dorche Isfahan and minimum rate of transmission achieved in white of Neishabour, so researchers suggest cultivate this genotype. Researchers suggest do more experiment for diagnose of resistant cultivar of onion, diagnose of ornamental hosts, trips and herbs that can transmit virus for provide control methods. Also, researchers suggest do more experiments for diagnose of other hosts of virus among of ornamental plants, crop plants and herbal plants.

ACKNOWLEDGEMENTS

Researchers thank Assist. Prof. Dr., from the Department of Plant Pathology, Faculty of Agriculture, Ferdowsi University of Mashhad (Iran). Researchers thank for editing the manuscript.

REFERENCES

- Bulajic, A., I. Djekic, J. Jovic, S. Krnjajic, A. Vucurovic and B. Krstic, 2009. Incidence and distribution of *iris yellow spot virus* on onion in serbia. *Plant Dis.*, 93: 976-982.
- Chatzivassiliou, E.K., I. Livieratos, G. Jenser and N.I. Katis, 2000. Ornamental plants and thrips populations associated with tomato spotted wilt virus in Greece. *Phytoparasitica*, 28: 257-264.
- Clark, M.F. and A.N. Adams, 1977. Characteristics of the microplate method of enzyme-linked immunosorbent assay for the detection of plant viruses. *J. Gen. Virol.*, 34: 475-483.
- Cortez, I., I.C. Livieratos, A. Derks, D. Peters and R. Kormelink, 1998. Molecular and serological characterization of *iris yellow spot virus*, a new and distinct tospovirus species. *Phytopathology*, 88: 1276-1282.
- Crowe, F.J. and H.R. Pappu, 2005. *Iris yellow spot virus*. *Plant Dis.*, 89: 105-105.
- Evans, C.K., S. Bag, E. Frank, J.R. Reeve, C. Ransom, D. Drost and H.R. Pappu, 2009. Natural infection of *Iris yellow spot virus* in Twoscale saltbush (*Atriplex micrantha*) growing in Utah. *Plant Dis.*, 93: 430-430.
- Gawande, S.J., A. Khar and K.E. Lawande, 2010. First Report of *Iris yellow spot virus* on garlic in India. *Plant Dis.*, 94: 1066-1066.

- Gent, D.H., L.J. Du Toit, S.F. Fichtner, S.K. Mohan, H.R. Pappu, and H.F. Schwartz, 2006. *Iris yellow spot virus*: An emerging threat to onion bulb and seed production. *Plant Dis.*, 90: 1468-1480.
- Gera, A., J. Cohen, R. Salomon and B. Raccach, 1998. *Iris yellow spot* tospovirus detected in onion (*Allium cepa*) in Israel. *Plant Dis.*, 82: 127-127.
- Ghotbi, T., N. Shahraeen and S. Winter, 2005. Occurrence of tospoviruses in ornamental and weed species in Markazi and Tehran provinces in Iran. *Plant Dis.*, 89: 425-429.
- Hsu, C.L., C.A. Hoepting, M. Fuchs, A.M. Shelton and B.A. Nault, 2010. Temporal dynamics of iris yellow spot virus and its vector, *Thrips tabaci* (Thysanoptera: Thripidae), in seeded and transplanted onion fields. *Environ. Entomol.*, 39: 266-277.
- Kritzman, A., H. Beckelman, S. Alexandrov, J. Cohen and M. Lampel *et al.*, 2000. Lisianthus leaf necrosis: A new disease of lisianthus caused by *Iris yellow spot virus*. *Plant Dis.*, 84: 1185-1189.
- Mandal, B., H.R. Pappu and A.K. Culbreath, 2001. Factors affecting mechanical transmission of *Tomato spotted wilt virus* to peanut (*Arachis hypogaea*). *Plant Dis.*, 85: 1259-1263.
- Mohan, S.K. and D.O. Wilson, Jr., 1989. Scape blight of onion. Proceedings of the National Onion Research Conference, August 16-18, 1989, Boise, ID., USA., pp: 103-104.
- Mumford, R.A., R. Glover, M. Daly, T. Nixon, V. Harju and A. Skelton, 2008. *Iris Yellow Spot Virus* (IYSV) infecting Lisianthus (*Eustoma grandiflorum*) in the UK: First finding and detection by real-time PCR. *Plant pathol.*, 57: 768-768.
- Pappu, H.R., I.M. Rosales and K.L. Druffel, 2008. Serological and molecular assays for rapid and sensitive detection of *Iris yellow spot virus* infection of bulb and seed onion crops. *Plant Dis.*, 92: 588-594.
- Pourian, H.R., M. Mirab-balou, M. Alizadeh and S. Orosz, 2009. Study on biology of onion thrips, *Thrips tabaci* (Thysanoptera, Thripidae) on cucumber (var. sultan) in laboratory conditions. *J. Plant Prot. Res.*, 49: 390-394.
- Pozzer, L., T. Nagata, M.I. Lima, E.W. Kitajima, R. de O. Resende, and A.C. de Avila, 1994. "Sapeca": An onion disease in Sub-Medio Sao Francisco region, Brazil, is caused by a tospovirus with a serologically distinct nucleocapsid protein. *Fitopatol. Bras.*, 19: 321-321.
- Pozzer, L., I.C. Bezerra, R. Kormelink, M. Prins, D. Peters, R.O. de Resende and A.C. de Avila, 1999. Characterization of a tospovirus isolate of *Iris yellow spot virus* associated with adisease in onion fields in Brazil. *Plant Dis.*, 83: 345-350.
- Silva, M.S., C.R. Martins, I.C. Bezerra, T. Nagata, A.C. de Avila and R.O. Resende, 2001. Sequence diversity of NS(M) movement protein of tospoviruses. *Arch. Virol.*, 146: 1267-1281.
- Smith, T.N., S.J. Wylie, B.A. Coutts and R.A.C. Jones, 2006. Localized distribution of *Iris yellow spot virus* within leeks and its reliable large-scale detection. *Plant Dis.*, 90: 729-733.
- Ward, L.I., Z. Perez-Egusquiza, J.D. Fletcher, F.M. Ochoa-Corona and J.Z. Tang *et al.*, 2008. First report of *Iris yellow spot virus* on *Allium cepa* in New Zealand. *Plant Pathol.*, 52: 406-406.