

## Relevance of Type of Breeding and Single Versus Double Sponge Inserts Followed by PMSG on Herri Ewes Estrus Exhibition and Pregnancy Rate

Mohamed Ali

Department of Animal Production, Qassim University, Qassim State, Saudi Arabia

**Abstract:** This study was designed to investigate effects of natural mating versus artificial insemination in ewes synchronized for estrus/ovulation by single versus double sponge inserts combined with PMSG. A 120 cyclic Herri ewes were utilized in the study. The 10 ewes served as control (C), given no treatment, 58 ewes were designed to be naturally mated at estrus exhibition, 47 of which were inserted with a single sponge impregnated in Fluoro Gestone Acetate (FGA) for 12 days and given 500 IU PMSG at sponge removal (SSNM-G1) and 11 ewes were given Double Sponges (DSNM-G2), the 1st sponge lasted 7 days, removed and replaced by a 2nd sponge for 5 days and at the 2nd sponge removal 500 IU PMSG was given (i.m). All ewes were visually observed for estrus signs and at onset of estrus, fertile rams were allowed to mate ewes. Time from sponge removal until estrus exhibition was recorded for each group. The artificially inseminated ewes were randomly allocated into 2 group, SSAI (G3, n = 31 ewes) and DSAI (G4, n = 21 ewes) in which ewes were given a single and double sponge inserts, respectively. Animals in G3 and G4 were timed inseminated with chilled fertile semen 48-50 h after sponge removal. All ewes were diagnosed for pregnancy 30 days post breeding by ultrasound. Proportion of ewes observed in estrus was highest in DSNM ewes (100%), followed by SSNM (91.5%) compared with C ewes (70%). Mean time interval from sponge removal till estrus was shorter in DSNM (38.1 h) than in SSNM (46.3 h). Likewise, conception and pregnancy rates were higher in DSNM (81.8 and 81.8%, respectively) than in SSNM (74.4 and 68%, respectively). Artificial insemination resulted in lower pregnancy rate (42.3%) with 35.4% in SSAI and 52.3% in DSAI ewes. Mean pregnancy rate in control ewes was 50%. In conclusion, inserting 2 sponges throughout 12 days combined with 500 IU PMSG at sponge removal shortened the interval from sponge withdrawal till exhibiting estrus, increased conception rate and pregnancy rate in Herri ewes.

**Key words:** Ewes, estrous synchronization, FGA, pregnancy rate, sponge

---

### INTRODUCTION

Improvement in fertility must first come from improved breeding management and only then from the use of Assisted Reproductive Technology (ART) such as estrus synchronization. Estrus synchronization facilitates the use of genetically superior sires through AI. It may, also enhance reproductive efficiency by shortening the breeding and lambing seasons. Synchronization of estrus involves the manipulation of the estrous cycle in cycling ewes by induction of estrus in anoestrus ewes, so that a large percentage of females come into estrus at a predetermined time. This procedure is known as timed breeding, breeding by appointment or mass mating (Olivera-Muzante *et al.*, 2013). For a successful estrous synchronization, the ewes should be closely synchronized with a rapid decline in circulating progesterone concentration, as well as synchronous growth and ovulation of a viable follicle.

There are 2 principles of controlling estrus and ovulation in ewe. The 1st principle is to prolong the life span of the CL thus delaying estrus. This can be attainable by administering progestin, such as a sponge or Controlled Internal Drug Release (CIDR) that mimics the function of the CL. The 2nd principle is to shorten the life of the CL has tening the onset of estrus. This is attainable by administering exogenous luteolytic agents such as prostaglandin  $F_{2\alpha}$ . However,  $PGF_{2\alpha}$  is effective only when a fully developed CL is present approximately between D5 and D7 of the estrous cycle (Hafez and Hafez, 2000).

Estradiol has been demonstrated to be a good marker for follicular function (Gonzalez-Bulnes *et al.*, 2005) for both sheep and cattle. In cattle, several studies have shown that healthy antral follicles contain high estradiol concentrations (Ireland and Roche, 1983) while atretic follicles contain low estradiol concentrations (Kruip and Dieleman, 1982) and are less able to release estradiol (Rouillier *et al.*, 1996). The lower values in plasma

estradiol concentration found in ewes treated with progestagens would indicate that follicular function might be altered in some of them. The basis for the administration of progestagens is the decrease in LH secretion (Goodman and Karsch, 1980) that prevents the occurrence of estrus, LH surge and ovulation until the withdrawal of the sponge. Maintenance of high levels of progestagen, also affects the expression of dominance effects from large follicles and increases follicular turnover (Leyva *et al.*, 1998). But, sometimes the release of progestagen from the sponges is too low at the end of the treatment; LH is not suppressed to the required level (Kojima *et al.*, 1992) and led to inadequate follicular development with persistent large follicles (Flynn *et al.*, 2000) that will affect fertility if ovulate (Vinoles *et al.*, 2001). The hypothesis was to evaluate the usefulness of inserting a second sponge within the 12 days combined with 500 IU PMSG at sponge removal on the enhancement of estrus occurrence and subsequent fertility and pregnancy rate in ewes.

## MATERIALS AND METHODS

A total of 120 cyclic Herri ewes,  $3\pm 5$  years of age and 40±45 kg body weight were used in this study. The experiment was conducted during Autumn 2010 at the Agriculture and Veterinary Research Center, Qassim University. The animals were managed under the same conditions on one farm. They were kept under semi-shaded yards, offered barley, alfalfa hay, clean water and integrated mineral licks. Animals were randomly allocated into 3 experiments. Experiment 1 (n = 10) ewes served as control, experiment 2 ewes (n = 58) served, as the naturally mated and subdivided into 2 groups, the Single Sponge Naturally Mated (SSNM) ewes (n = 47) and the Double Sponge Naturally Mated (DSNM) ewes (n = 11). However, ewes were designated for AI and subdivided into 2 groups, the Single Sponge AI (SSAI, n = 31) and the Double Sponge AI ewes (DSAI, n = 21). Figure 1 illustrated the design and the time schedule for various animal treatments.

**Experiment 1 (control):** The 1 ewes (n = 10) served, as control receiving no vaginal pessaries and introduced to the males at the same time when ewes of experiment 2 were introduced to the males.

**Experiment 2 (natural mating):** Ewes (n = 58) were randomly allocated into the following groups; G1 (SSNM, n = 47). Ewes were vaginally inserted with a single sponges impregnated with 40 mg of Flouro Gestone Acetate (FGA) and left for 12 days. Group 2 (DSNM, n = 11): Ewes were inserted with a single vaginal sponges




Fig. 1: Sponge-PMSG protocols in experiment 1 and 2 In G2 and G4 sponge was inserted for 12 days and PMSG injection at sponge removal. In G3 and G5, ewes were treated with single vaginal sponges for 7 days and then replaced with new vaginal sponge for the rest time (5 days) and PMSG injection at sponge removal. There was no treatment in the experiment 1

for 7 days and then replaced with a new vaginal sponge for the rest 5 days. At sponge removal, all ewes were injected (i.m) with 500 IU of Pregnant Mare Serum Gonadotropine (PMSG). At the onset of estrus, natural mating was carried out by the use of fertile rams in a ratio of 1:5. The males were introduced in the herd after pessary removal for a total period of 96 h. The duration from sponge removal until estrus exhibition was recorded. After 30 days of natural mating, pregnancy was confirmed by ultrasound (Esaote, pie medical, Netherlands).

**Experiment 3 (artificial insemination):** Females (n = 52) were randomly allocated into 2 groups; group 3 (SSAI, n = 31), ewes were inserted with a single sponge for 12 days. While group 4 (DSAI, n = 21), ewes were inserted with a sponge 7 days and then replaced by another sponge for the remained 5 days. At sponge removal, all ewes were injected (i.m) with 500 IU PMSG. Semen from a fertile ram was collected using an artificial vagina. Having determined seminal characteristics, it was

diluted with egg yolk-glucose-citrate extender. All ewes were artificially inseminated with fresh diluted semen containing a dose of 200 millions total sperm. Cervical artificial insemination was performed 48-50 h after sponge removal (Timed AI). After 30 days of artificial insemination, pregnancy was diagnosed by ultrasound.

**Statistical analysis:** Data were analysed using a statistical software SPSS release 17.0. Reproductive parameters among 5 group treatments were compared by ANOVA and Chi-square. Data are presented as arithmetic mean $\pm$ SEM. Significance was considered at  $p < 0.05$ .

## RESULTS AND DISCUSSION

Table 1 shows the effect of synchronization treatment on estrus response in the Herri ewes. In the experiment 2 (natural mating), a total of 54 out of 58 (93.1%) ewes exhibited estrus. The percentage of ewes observed in estrus was highest in the DSNM (100%), however only 91.5% of SSNM ewes exhibited estrus. Although, the difference was not statistically significant ( $p > 0.05$ ). Ewes given the second sponge insert have shown estrus shortly (after 38.1 h) after sponge removal as compared with those given a sole sponge (46.3 h). In case of single sponge insert, the estrus exhibition duration scattered between 24-144 h, however inserting a second sponge intensified the estrus duration within short time ( $< 24$ -48 h). The pregnancy rate after synchronization treatment in the G1 and G2 were 68.0 and 81.8%, respectively which were higher than control (50%). Despite this enhancement, there was no statistical difference. In experiment 3 (AI), likewise, the second sponge insert tended to increase pregnancy rate (52.3%) than the single sponge artificially inseminated ewes (35.4%). Comparing natural mating with artificial insemination on the pregnancy rates was in a favor of natural mating (70.6%) which was higher than AI (42.3%).

Distribution of ewes exhibiting estrus over time following sponge removal in experiment 2 is shown in Fig. 2. The 40 out of 43 ewes (93%) exhibited estrus within 72 h of sponge removal in the SSNM ewes, however in ewes given the second sponge 100% exhibited estrus within 48 h of sponge removal. No estrus observation was conducted for the AI ewes, since they were designed to be artificially inseminated at predetermined time (48 h after sponge removal). Control ewes exhibited estrus in only 7 out of 10 ewes (70%) with wider time frame of 144 h.

Insertion of a 2nd progesterone sponge not only accelerated the onset of estrus but it also increased the proportion of estrus occurrence in synchronized ewes. The 2nd dose of progesterone might optimized the internal progesterone concentration leading to an optimum peak. When the 2nd sponge insert was removed a sharp decline in peripheral progesterone has been occurred resulting in a better trigger of LH peak. The resultant LH peak might caused better ovulation, resulting in improved pregnancy rate (81.8%). The use of PMSG in conjunction with intravaginal progestagen treatment was found to efficient for estrus induction synchronization and ovulation (Ainsworth *et al.*, 1983; Barrett *et al.*, 2004;




Fig. 2: Distribution of ewes exhibiting estrus over time following synchronization treatment in experiment 2; interval (h) between sponage removal and estrus exhibition

Table 1: Estrus response and pregnancy rate following synchronization treatment in the Herri ewe using natural mating and artificial insemination

Groups	Number	No. of ewes in estrus (%)	Sponge removal to estrus (h)	Conception rate (%)	No. of pregnant ewe (%)
<b>Experiment 1</b>					
Control	10	7 (70.0)	52.0 $\pm$ 15.8	71.4	5 (50.0)
<b>Experiment 2 (NM)</b>					
G1 (SSNM)	47	43 (91.5) <sup>a</sup>	46.3 $\pm$ 3.3 <sup>b</sup>	74.4 <sup>a</sup>	32 (68.0) <sup>aA</sup>
G2 (DSNM)	11	11 (100) <sup>a</sup>	38.1 $\pm$ 1.4 <sup>a</sup>	81.8 <sup>a</sup>	9 (81.8) <sup>aA</sup>
Total	58	54 (93.1)	44.05 $\pm$ 2.2	75.9	41 (70.6) <sup>A</sup>
<b>Experiment 3 (AI)</b>					
G3 (SSAI)	31	-	-	-	11 (35.4) <sup>aB</sup>
G4 (DSAI)	21	-	-	-	11 (52.3) <sup>aA</sup>
Total	52	-	-	-	22 (42.3) <sup>B</sup>

<sup>a,b</sup> Values with different superscripts in the same column and experiment differ significantly at  $p < 0.05$ ,  $\chi^2$  and ANOVA; <sup>A,B</sup> Values with different superscripts in the same column between experiments differ significantly at  $p < 0.05$ ,  $\chi^2$

Koyuncu and Alticekic (2010) in ewes. Similar finding was reported by Dogan and Nur (2006) confirming that the double sponge inserts within 14 days accompanied by the injection of PMSG at sponge removal shortened the time duration till estrus, than in the single-sponge treated ewes in the non-breeding season.

Concomitantly, other researchers (Ungerfeld and Rubianes, 1999; Das *et al.*, 2000; Simonetti *et al.*, 2000; Vinales *et al.*, 2001; Dogan and Nur, 2006) found similar interval when used double sponge impregnated with progesterone analogs. When ewes were given progesterone (Das *et al.*, 2000) or FGA or MAP intravaginal sponges (Ungerfeld and Rubianes, 1999; Simonetti *et al.*, 2000) alone or combined with PMSG (Greyling *et al.*, 1997; Zarkawi *et al.*, 1999; Vinales *et al.*, 2001; Ungerfeld and Rubianes, 2002; Zeleke *et al.*, 2005) the estrus response was recorded until 120 h after pessary removal.

These results showed that the time interval to onset estrus in the control and SSNM ewes ranged between 24-144 h whereas in DSNM ewes of this interval has been shortened (25-48 h). Therefore, ewes treated with one sponge tended to have longer intervals to estrus than those treated with two sponges. Researchers have reported the onset of estrous to occur within 24-144 h following progestagen or progesterone withdrawal (Ungerfeld and Rubianes, 1999; Das *et al.*, 2000; Simonetti *et al.*, 2000; Vinales *et al.*, 2001). Ewes came into heat between 18 and 96 h after sponge withdrawal with the highest incidence of estrous onset occurring between 30 and 60 h. The distribution of estrous in the present study ewes was similar to that reported elsewhere (Zarkawi *et al.*, 1999; Ungerfeld and Rubianes, 1999; Simonetti *et al.*, 2000; Dogan and Nur, 2006).

Additionally, the findings of this study demonstrated some ewes failed to display estrus after estrus synchronization treatment and that was higher in SSNM and control ewes. These findings indicate that there is great variation in the estrus response in Herri ewes after sponge treatment. This variation could be due to various factors and conditions that occur before or during the estrus synchronization treatment. Dogan and Nur (2006) demonstrated that 4 out of 13 ewes did not show any overt signs of estrous but were diagnosed pregnant at day 54 after AI. Such a finding is in agreement with a previous report in Boer and indigenous goats (Greyling and Van der Nest, 2000). Allison and Robinson (1970) suggested that these silent ovulations may be related to inadequate endogenous progesterone levels. Besides, absence of estrus and ovulation may be due to insufficient gonadotrophic hormone released by the pituitary to a poor response by ovary to the exogenous

PMSG or variation in responsiveness of animals to PMSG (Akusu and Egbunike, 1984). Moreover, in the current study only two visual observation periods (07:00-09:00 and 18:00-20:00 h) were applied, this time schedule might was not enough for visualizing those ewes exhibiting short estrus signs in the rest of the day (24 h). Failure of estrus detection might be attributed to an inefficient detection scheme and/or the occurrence of true silent ovulations that make it difficult to determine patterns of cyclicity (Garcia, 1990).

Ewes given Double Sponge (DS) either naturally mated or artificially inseminated gave better pregnancy rates than their counterparts give Single Sponge (SS). It was reported that using one sponge can lead to a decrease in the conception rate after the treatment. The events that bring about ovulation are highly dependent and integrated. Slight changes may disturb the maturation of oocytes and timing of ovulation which result in impaired developmental competence of oocytes leading to an early embryonic death (Bage *et al.*, 2000). In addition with these changes in timing of ovulation, the chance of normal AI timing is impaired, resulting in fertilization failure and consequently less pregnancy (Bage *et al.*, 2000). During this period, the follicular wave pattern was not maintained and a large follicle persisted following spontaneous luteolysis (Larson and Ball, 1992). The sequential relationship of the low progesterone increased the frequency of pulses of LH, a persistent largest follicle increased secretion of estradiol-17  $\beta$  and reduced fertility is widely accepted as one of the causes and effects (Inskeep, 2004). From these aged large follicles, abnormal ova will ovulate and this may be one of the reasons for low pregnancy rates (Inskeep, 2004). This could also be the reason as to why we observed low pregnancy rate in the SS ewes in the present study. One other critical factor that reduced sperm transport can occur after long period of low progesterone treatment (Larson and Ball, 1992). In another study by Shaham-Albalancy *et al.* (1997), they reported that concentrations of progesterone before estrus altered endometrial morphology during the subsequent estrous cycle. Low concentrations of progesterone (2.1-2.3 ng mL<sup>-1</sup>) during that period increased subsequent secretion of PGF<sub>2 $\alpha$</sub>  in response to oxytocin, as measured by its major metabolite (Shaham-Albalancy *et al.*, 2001). These effects might lead to lowered embryonic survival and also a decrease in fertility. Thus, low pregnancy rate in single sponge ewes might began very early due to the exposure to lowered progesterone level that may lead to embryonic death.

The physiological explanation of the lower pregnancy rates in AI versus natural mating is due to the predetermined timing of inseminating the ewes designed for AIs. As it is clear that estrus/ovulation occurrence scattered over 144 h in SS ewes, this would impaired the

fertilization process in a specific percentage of ewes (Pearce and Robinson, 1985; Baril *et al.*, 1993). However, in double sponge ewes the 52.3% of pregnancy rate resulting from artificial insemination would be acceptable under the experiment condition. It still comparable to the percentage obtained in SSNM ewes.

## CONCLUSION

Vaginally insertion of 2 progestagen sponge in Herri ewes resulted in a shorter intensifying interval from the sponge with drawal to estrus. This regime increased the possibility of ova fertilization and subsequent pregnancy rate. Further, studies warrant to be conducted on large-scale number of ewes from various local breeds inside and outside the breeding season.

## ACKNOWLEDGEMENT

I wish to express my sincere thanks and deepest gratitude to Prof. Dr. Muostafa Zeitoun for editing my study.

## REFERENCES

- Ainsworth, L., R. Lachance and F. Labrie, 1983. Effects of progestagen treatments and PMSG on the induction of the preovulatory LH discharge in ewes. *Anim. Reprod. Sci.*, 5: 281-286.
- Akusu, M.O. and G.N. Egbunike, 1984. Fertility of the West African Dwarf goat in its native environment following prostaglandin  $F_2\alpha$  induced estrus. *Vet. Q.*, 6: 173-176.
- Allison, A.J. and T.J. Robinson, 1970. The effect of dose level of intravaginal progestagen on sperm transport, fertilization and lambing in the cyclic merino ewe. *J. Reprod. Fertil.*, 22: 515-531.
- Bage, R., M. Forsberg, H. Gustafsson, B. Larsson and H. Rodriguez-Martinez, 2000. Effect of ACTH-challenge on progesterone and cortisol levels in ovariectomised repeat breeder heifers. *Anim. Reprod. Sci.*, 63: 65-76.
- Baril, G. B. Leboeuf and J. Saumande, 1993. Synchronization of estrus in goats: The relationship between time of occurrence of estrus and fertility following artificial insemination. *Theriogenology*, 40: 621-628.
- Barrett, D.M., P.M. Bartlewski, M. Batista-Arteaga, A. Symington and N.C. Rawlings, 2004. Ultrasound and endocrine evaluation of the ovarian response to a single dose of 500 IU of eCG following a 12-day treatment with progestogen-releasing intravaginal sponges in the breeding and nonbreeding seasons in ewes. *Theriogenology*, 61: 311-327.
- Das, G.K., S.M.K. Naqvi, R. Gulyani, S.R. Pareek and J.P. Mittal, 2000. Effect of two doses of progesterone on estrus response and fertility in acyclic crossbred Bharat Merino ewes in a semi-arid tropical environment. *Small Rumin. Res.*, 37: 159-163.
- Dogan, I. and Z. Nur, 2006. Different estrous induction methods during the non-breeding season in Kivircik ewes. *Vet. Med.*, 51: 133-138.
- Flynn, J.D., P. Duffy, M.P. Boland and A.C.O. Evans, 2000. Progestagen synchronisation in the absence of a corpus luteum results in the ovulation of a persistent follicle in cyclic ewe lambs. *Anim. Reprod. Sci.*, 62: 285-296.
- Garcia, M., 1990. Milk progesterone profiles in crossbred Brown Swiss  $\times$  Nellore cattle following natural service. *Theriogenology*, 33: 1105-1111.
- Gonzalez-Bulnes, A., P. Veiga-Lopez, P. Garcia, R.M. Garcia-Garcia and C. Ariznavarreta *et al.*, 2005. Effects of progestagens and prostaglandin analogues on ovarian function and embryo viability in sheep. *Theriogenology*, 63: 2523-2534.
- Goodman, R.L. and F.J. Karsch, 1980. Pulsatile secretion of luteinizing hormone: Differential suppression by ovarian steroids. *Endocrinology*, 107: 1286-1290.
- Greyling, J.P.C., J.A. Erasmus, G.J. Taylor, S. Van der Merwe, 1997. Synchronization of estrus in sheep using progestagen and inseminating with chilled semen during the breeding season. *Small Rumin. Res.*, 26: 137-143.
- Greyling, J.P.C. and M. Van der Nest, 2000. Synchronization of oestrus in goats: Dose effect of progestagen. *Small Rumin. Res.*, 36: 201-207.
- Hafez, B. and E.S.E. Hafez, 2000. Reproduction in Farm Animals. Williams and Wilkins, Lippincott, Pages: 512.
- Inskip, E.K., 2004. Preovulatory, postovulatory and postmaternal recognition effects of concentrations of progesterone on embryonic survival in the cow. *J. Anim. Sci.*, 82: 24-39.
- Ireland, J.J. and J.F. Roche, 1983. Development of nonovulatory antral follicles in heifers: Changes in steroids in follicular fluid and receptors for gonadotropins. *Endocrinology*, 112: 150-156.
- Kojima, N., T.T. Stumpf, A.S. Cupp, L.A. Werth and M.S. Roberson *et al.*, 1992. Exogenous progesterone and progestins as used in estrous synchrony regimens do not mimic the corpus luteum in regulation of luteinizing hormone and 17  $\beta$ -estradiol in circulation of cows. *Biol. Reprod.*, 47: 1009-1017.
- Koyuncu, M. and S.O. Alticekic, 2010. Effects of progestagen and PMSG on estrous synchronization and fertility in Kivircik ewes during natural breeding season. *Asian-Aust. J. Anim. Sci.*, 23: 308-311.

- Kruij, T.A.M. and S.J. Dieleman, 1982. Macroscopic classification of bovine follicles and its validation by micromorphological and steroid biochemical procedures. *Reprod. Nutr. Dev.*, 22: 465-473.
- Larson, L.L. and P.J.H. Ball, 1992. Regulation of estrous cycles in dairy cattle: A review. *Theriogenology*, 38: 255-267.
- Leyva, V., B.C. Buckrell and J.S. Walton, 1998. Regulation of follicular activity and ovulation in ewes by exogenous progestagen. *Theriogenology*, 50: 395-416.
- Olivera-Muzante, J., J. Gil, C. Vinales and S. Fierro, 2013. Reproductive outcome with GnRH inclusion at 24 or 36 h following a prostaglandin F2 $\alpha$ -based protocol for timed AI in ewes. *Anim. Reprod. Sci.*, 138: 175-179.
- Pearce, D.T. and T.J. Robinson, 1985. Plasma progesterone concentrations, ovarian and endocrinological responses and sperm transport in ewes with synchronized oestrus. *J. Soc. Reprod. Fertil.*, 75: 49-62.
- Rouillier, P., L.A. Guilbault, J.G. Lussier and P. Matton, 1996. Changes in morphological appearance and functional capacity of recruited follicles in cows treated with FSH in the presence or absence of a dominant follicle. *Theriogenology*, 46: 1053-1061.
- Shaham-Albalancy, A., A. Nyska, M. Kaim, M. Rosenberg, Y. Folman and D. Wolfenson, 1997. Delayed effect of progesterone on endometrial morphology in dairy cows. *Anim. Reprod. Sci.*, 48: 159-174.
- Shaham-Albalancy, A., Y. Folman, M. Kaim, M. Rosenberg and D. Wolfenson, 2001. Delayed effect of low progesterone concentrations on bovine uterine PGF(2 $\alpha$ ) secretion in the subsequent oestrous cycle. *Reproduction*, 122: 643-648.
- Simonetti, L., M.R. Blanco and J.C. Gardon, 2000. Estrus synchronization in ewes treated with sponges impregnated with different doses of medroxyprogesterone acetate. *Small Rumin. Res.*, 38: 243-247.
- Ungerfeld, R. and E. Rubianes, 1999. Estrus response to the ram effect in Corriedale ewes primed with medroxyprogesterone during the breeding season. *Small Rumin. Res.*, 32: 89-91.
- Ungerfeld, R. and E. Rubianes, 2002. Short term primings with different progestogen intravaginal devices (MAP, FGA and CIDR) for eCG-estrous induction in anestrus ewes. *Small Rumin. Res.*, 46: 63-66.
- Vinales, C., M. Forsberg, G. Banchemo and E. Rubianes, 2001. Effect of long-term and short-term progestagen treatment on follicular development and pregnancy rate in cyclic ewes. *Theriogenology*, 55: 993-1004.
- Zarkawi, M., M.R. Ai-Merestani and M.F. Wardeh, 1999. Induction of synchronized oestrous and early pregnancy diagnosis in Syrian Awassi ewes, outside the breeding season. *Small Rumin. Res.*, 33: 99-102.
- Zelege, M., J.P.C. Greyling, L.M.J. Schwalbach, T. Muller and J.A. Erasmus, 2005. Effect of progestagen and PMSG on oestrous synchronization and fertility in dorper ewes during the transition period. *Small Rumin. Res.*, 56: 47-53.