

## On the Use of Orthogonality Relations to Construct Character Tables of Finite Group

H.S. Ndakwo

Department of Mathematical Sciences  
Nasarawa State University, Keffi, Nasarawa State Nigeria

**Abstract:** Group characters are very fundamental in the study of group representation. In this paper we employed the orthogonality relation in constructing the character table which shows the difference irreducible characters for each conjugacy class. The first kind is used along the rows, while the second kind is used along the columns. Relevant details of this method are described following.

**Key words:** Group characters, representation, orthogonality relation, irreducible, conjugacy class, inner product.

### INTRODUCTION

The complete information about the character of  $G$  is conveniently displayed by Isaacs, (1956) in a character table which lists the values of the  $K$  simple character for all elements. We know that a character is constant on each of the conjugacy classes  $C_a$ . If  $x \in C_a$ , we put  $\chi(x) = \chi^{(a)}$ . Thus it is sufficient to record the values  $\chi^{(a)}$ ,  $1 \leq a \leq k$

Denoting the number of elements in  $C_a$  by  $h_k^{(a)}$ , we have the class equation

$$h^{(1)} + h^{(2)} + \dots + h^{(k)} = g \quad (1)$$

Unless the contrary is stated. We adhere to the convention that  $G = \{1\}$  and that  $F$  is the trivial representation  $F_1(x) = 1, x \in G$

The Table 1 below represent a typical character table, the body of the table is a  $K$  square matrix whose rows correspond to the different characters, while each column contain the values of all simple character for a particular conjugacy class.

**Definition 1:** A generalized character of a group  $G$  is a function of the form

$$\sum_{i=1}^s a_i \chi_i$$

	$C^1$	$C^2 \dots$	$C^A \dots$	$C^K$
$\chi^1$	$h^{(1)}$	$h^{(2)} \dots$	$h^{(a)} \dots$	$h^0$
$\chi^2$	$f^1$	$1 \dots$	$1 \dots$	$1$
$\vdots$	$\vdots$	$\chi_2^{(2)} \dots$	$\chi_2^{(a)} \dots$	$\chi_2^{(k)}$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$\chi^1$	$f_i$	$\chi_i^{(2)}$	$\chi_k^{(a)}$	$\chi_k^{(k)}$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$\chi^k$	$f_k$	$\chi_k^{(2)} \dots$	$\chi_k^{(a)} \dots$	$\chi_k^{(k)}$

where  $\chi_i$  are irreducible and  $a_i \in \mathbb{Z}$ . Generalized characters are class functions that are constant on any conjugacy class.

**Definition 2:** Characters of degree one are called linear characters by Feit (1971). In particular, the function with constant value 1 on  $G$  is a linear  $F$ -character. It is called the principal  $F$ -character.

**Definition 3:** Let  $\chi$  be an  $F$ -representation of  $G$ . Then the  $F$ -character  $\chi$  of  $g$  afforded by  $x$  is a function given by  $\chi(g) = \text{tr}(g)$ .

### Lemma 4

- Similar  $F$ -representation of  $G$  afforded equal characters.
- Characters are constant on the conjugacy classes of a group.

### Proof

- If  $P$  is non singular then  $\text{tr}(P^{-1}AP) = \text{tr}(PP^{-1}A) = \text{tr}(A)$ . Both (a) and (b) follow from this observations.
- Observe that  $\chi(h^{-1}gh) = \chi(h)^{-1} \chi(g) \chi(h)$  if  $\chi$  is a representation of  $G$  and hence  $\text{tr}(\chi(h^{-1}gh)) = \text{tr}(\chi(g))$ . Hence the result.

### Corollary 5

The group  $G$  is abelian if and only if every irreducible character is linear.

### Proof

Let  $k$  be the number of classes of  $G$ . Then  $K = |G|$  if and only if  $G$  is abelian.  
Now

	S <sup>3</sup>		
	C <sub>1</sub>	C <sub>2</sub>	C <sub>3</sub>
h <sup>(a)</sup>	1	3	2
χ <sub>1</sub>	1	1	1
χ <sub>2</sub>	1	-1	1
χ <sub>3</sub>	2	0	-1

$$|G| = \sum_{i=1}^k \chi_i(1)^2 \text{ and } \chi_i(1) \geq 1$$

and for all i it follows that  $K = |G|$  if and only if  $\chi_i(1) = 1$  for all i. Hence the result.

#### NOTE

For a particular group G, the irreducible characters are usually presented in a character table. A square array of complex numbers whose rows correspond to the  $\chi_i$  and whose columns correspond to the classes  $C^1$ .

**Example 6:** An example of a character table is the accompanying one for the symmetric groups  $S^3$  on three symbols. It has three conjugacy classes namely;

$$\begin{aligned} C^1 &= \{1\} \\ C^2 &= \{(12), (13), (23)\} \\ C^3 &= \{(123), (132)\} \end{aligned}$$

Therefore,  $S^3$  has three irreducible representations. The character Table 2 of  $S^3$  is given below

This method for computing the coefficient is by using orthogonality relations which we are about to derive.

These relations are useful in the construction of character tables.

The key to the orthogonality relations is to compute explicitly the coefficients of the group elements in the  $C^1$ 's in terms of the characters.

#### Definition 7:

Let  $\phi$  and  $\psi$  be class functions on a group G. Then (Dornhoff, 1971)

$$[\phi, \psi] = \frac{1}{|G|} \sum_{g \in G} \phi(g) \psi(g)$$

is the inner product of  $\phi$  and  $\psi$ .

#### Properties of inner product 8

$$[\phi, \psi] = \overline{[\psi, \phi]}$$

$$[\phi, \phi] > 0 \text{ Unless } \phi \neq 0$$

$$[C_1 \phi_1 + C_2 \phi_2, \psi] = C_1 [\phi_1, \psi] + C_2 [\phi_2, \psi]$$

$$[\phi, C_1 \psi_1 + C_2 \psi_2] = \overline{C_1} [\phi, \psi_1] + \overline{C_2} [\phi, \psi_2]$$

#### Remark 9

The  $[\ ]$  has all the properties usually used to define an inner product in linear algebra and analysis. We know that  $\text{irr}(g)$  is a basis for the space of class functions and it is the content of the orthogonality relation which is in fact an orthonormal basis that is

$[\chi_i, \chi_j] = \delta_{ij}$ . This yields the promised method for expressing an arbitrary class functions in terms of the irreducible characters, for if,

$$[\phi, \chi_i] = C_i. \text{ Then}$$

$$\phi = \sum C_i \chi_i$$

#### Definition 10 (Alternative form of the inner product)

If X lies in  $C^a$ , we let

$$\phi(x) = \phi(a)$$

$$\psi(x) = \psi(a)$$

$$\psi(x^{-1}) = \overline{\psi(a)}$$

Then the inner product of  $\phi$  and  $\psi$  is

$$[\phi, \psi] = \frac{1}{g} \sum_{x \in g} \phi(x) \psi(x^{-1}) = \frac{1}{g} \sum_{a=1}^k h^{(a)} \phi(a) \overline{\psi(a)} \quad (2)$$

#### Note

If  $\chi_1$  and  $\chi_2$  are simple characters, the character relation of the first kind by (Ledermann, 1976) states that

$$\frac{1}{g} \sum_{a=1}^k h^{(a)} \chi_1^{(a)} \chi_2^{(a)} = \begin{cases} 1, & \chi_1 = \chi_2 \\ 0, & \chi_1 \neq \chi_2 \end{cases}$$

Applying the result to all the rows of the character Table 2 we have that

$$\frac{1}{g} \sum_{a=1}^k h^{(a)} \chi_1^{(a)} \chi_2^{(a)} = \delta_{ij} \quad (3)$$

Where  $i, j = 1, 2, \dots, K$ . We can express this information in a more compact form by introducing the  $k \times k$  matrix,

$$(u_{ia}) = \chi_i^{(a)} \sqrt{\frac{h^{(a)}}{g}} \quad (4)$$

Then Eq. (4) states that the rows of U form a unitary matrix that is

$$U U^{-1} = I \quad (5)$$

Where t denotes transposition. U and U<sup>t</sup> are inverse to each other and therefore commute. Hence

$$UU^t = I$$

And on taking the conjugate complex, we have This means that the columns of U also form a unitary orthogonal system, that is

$$UU^t = I$$

Or in terms of characters,

$$\sum_{i=1}^k U_{ia} \overline{U_{ib}} = \frac{g}{h^{(a)}} \delta_{ab} \quad (6)$$

$$\sum_{i=1}^k \chi_i^{(a)} \overline{\chi_i^{(b)}} = \frac{g}{h^{(a)}} \delta_{ab}$$

Equation (6) is called orthogonality relation of the second kind.

## MAIN RESULT

### THEOREM 11 (ORTHOGONALITY RELATIONS)

(i) If ij then

$$\sum_{g \in G} \chi_i(g) \chi_j(g^{-1}) = 0$$

Alternatively,

$$\sum_{k=1}^s h_k \chi_i(g) \chi_j(\chi_k^{-1}) = |G|, k \in G$$

$$(ii) \sum_{g \in G} \chi_i(g) \chi_i(g^{-1}) = |G|.$$

Alternatively

$$\sum_{k=1}^s h_k \chi_i(\chi) \chi_j(\chi_k^{-1}) = |G|$$

$$(iii) \sum_{k=1}^s \chi_k(\chi_i) \chi_k(\chi_j^{-1}) = 0$$

if i ≠ j

$$(iv) \sum_{k=1}^s \chi_k(\chi_i) \chi_k(\chi_i^{-1}) = \frac{|G|}{h_i} = |c(x_i)|$$

(v) The  $\chi_i$  are linearly independent over C as functions from the set of conjugacy class of C.

$$(vi) \sum_{k=1}^s \chi_k^2(1) = |G|$$

### Proof

We have

$$\chi_i(g) = \text{tr}(\psi_i(g)) = \text{tr}(a_{ij}(g)) = \sum_m a_{mm}(g)$$

and similarly,

$$\chi_j(g) = \sum_n b_{nn}(g).$$

$$\begin{aligned} \sum_{g \in G} \chi_i(g) \chi_j(g^{-1}) &= \sum_g \left( \sum_m a_{mm}(g) \right) \left( \sum_n b_{nn}(g^{-1}) \right) \\ &= \sum_{m,n} \sum_g a_{mm}(g) b_{nn}(g^{-1}) = 0 \end{aligned}$$

If i ≠ j (for  $V_i \cong V_j$ ). Hence (i)

$$\sum_m a_{mm}(g) \sum_n b_{nn}(g^{-1}) = 0$$

(ii). Unless m = n. The case where i = j and a = b we get.

$$\begin{aligned} \sum_m \sum_g a_{mm}(g) a_{mm}(g^{-1}) &= \sum_m \delta_{mm} \frac{|G|}{\dim V_i} = \\ \sum_m \frac{|G|}{\dim V_i} &= \dim V_i \frac{|G|}{\dim V_i} = |G| \end{aligned}$$

Hence (ii).

Let Y be the matrix with (i, j)  $\chi_j(x_i^{-1})$  entry. and  $\chi$  are the matrix with (i, j) entry  $h_i \chi_i(x_j)$ . But X and Y are square matrix. So XY = I  $\Rightarrow$  YX = I Hence (iii) and (iv)

(v) Is a consequence of (i) and (ii).

Suppose  $\chi_i$  were linear combination, say,

$$\begin{aligned} \chi_i &= a_2 \chi_2 + a_3 \chi_3 + \dots + a_n \chi_n \text{ then,} \\ \sum h_m \chi_i(x_m) \chi_j(x_m^{-1}) &= 0 \text{ for all} \\ \sum h_m \chi_i(x_m) \left( \sum_{j=2}^s a_j \chi_j(x_m^{-1}) \right) &= 0 \Rightarrow \\ \sum h_m \chi_i(x_m) \chi_j(x_m^{-1}) &= 0 \end{aligned}$$

But the left hand side is equal to |G| by (ii) and |G| ≠ 0. Thus a contradiction. (vi) Is a consequence of (iv) with

$$\begin{aligned} \chi_i &= 1 = \chi_i^{-1} \text{ but } \chi_i(1) = \dim V_i = n_i \text{ and we have} \\ h_1^2 + h_2^2 + \dots + h_n^2 &= |G| \end{aligned}$$

### Example 12 (Construction of character tables)

(i) Using the above theorem to construct the character table for symmetric group of degree 3.

	S <sup>3</sup>		
	C <sub>1</sub>	C <sub>2</sub>	C <sub>3</sub>
h <sup>(a)</sup>	1	3	2
χ <sub>1</sub>	1	1	1
χ <sub>2</sub>	a	c	d
χ <sub>3</sub>	b	e	f

The symmetric group is the group of all permutations of the. It has six elements and three conjugacy classes namely,

$$\begin{aligned} C_1 &= \{1\} \\ C_2 &= \{(12), (13), (23)\} \\ C_3 &= \{(123), (132)\} \end{aligned}$$

Therefore the symmetric group of degree three has 3 irreducible representations.

Let the values of χ<sub>2</sub> for the classes C<sub>1</sub>, C<sub>2</sub> and C<sub>3</sub> be a, c and d respectively and the values for χ<sub>3</sub> of the classes C<sub>1</sub>, C<sub>2</sub> and C<sub>3</sub> be b, e and f respectively.

Thus the character Table 3 or the symmertwric group of degree three is

Now to find a, b, c, d, e and f. However,

$$1^2 + a^2 + b^2 = 6 \Rightarrow a^2 + b^2 = 5$$

That is  $1^2 + 2^2 = 5 \Rightarrow a=1, b=2$ .

By using the orthogonality relations of the second kind and taking the inner products to find the remaining variables.

Now if a = 1, b = 2, we have

$$\begin{aligned} \chi_1^{(1)} \cdot \chi_1^{-(2)} + \chi_2^{(1)} \cdot \chi_2^{-(2)} + \chi_3^{(1)} \cdot \chi_3^{-(2)} &= 0 \\ 1.1 + 1.c + 2.e &= 0 \\ 1 + c + 2e &= 0 \end{aligned} \quad (7)$$

If a = 2, b = 2, we have

$$\begin{aligned} \chi_1^{(2)} \cdot \chi_1^{-(2)} + \chi_2^{(2)} \cdot \chi_2^{-(2)} + \chi_3^{(2)} \cdot \chi_3^{-(2)} &= \frac{6}{3} \\ 1.1 + c.c + e.e &= 2 \\ c^2 + e^2 &= 1 \end{aligned}$$

But c = -1-2e so (8) becomes

$$\begin{aligned} (-1 - 2e)^2 + e^2 &= 1 \\ 1 + 4e + 5e^2 &= 1 \\ 4e + 5e^2 &= 0 \end{aligned}$$

e = 0 or -4/5 put e = 0 in Eq. 7 we have c = -1. And if a = 1, b = 3 we have,

	S <sup>3</sup>		
	C <sub>1</sub>	C <sub>2</sub>	C <sub>3</sub>
h <sup>(a)</sup>	1	3	2
χ <sub>1</sub>	1	1	1
χ <sub>2</sub>	1	-1	1
χ <sub>3</sub>	2	0	-1

$$\begin{aligned} \chi_1^{(1)} \cdot \chi_1^{-(3)} + \chi_2^{(1)} \cdot \chi_2^{-(3)} + \chi_3^{(1)} \cdot \chi_3^{-(3)} &= 0 \\ 1.1 + 1.d + 2.f &= 0, \\ d &= -1 - 2f \text{ if} \end{aligned} \quad (9)$$

a = 3, b = 3, we have

$$\begin{aligned} \chi_1^{(3)} \cdot \chi_1^{-(3)} + \chi_2^{(3)} \cdot \chi_2^{-(3)} + \chi_3^{(3)} \cdot \chi_3^{-(3)} &= \frac{6}{2} \\ 1.1 + d^2 + f^2 &= 3 \Rightarrow d^2 + f^2 = 3 \\ \text{But } d &= -1 - 2f \\ (-1 - 2f)^2 + f^2 &= 2 \\ 5f^2 + 4f - 1 &= 0 \\ f &= -1 \text{ or } \frac{1}{5} \end{aligned}$$

Put f = -1 in Eq. (9) we have d = 1.

Therefore the required character Table 4 for the symmetric group of degree three is

### THEOREM 13

Let N be a normal subgroup of G and suppose that A<sub>0</sub>(N<sub>x</sub>) is a representation of degree m of the group, G/N then by (Curtis and Reiner (1962) we have the following A(x) = A<sub>0</sub>(N<sub>x</sub>), x ∈ G, defines a representation of G “lifted from G/N”.

If ϕ<sub>0</sub>(N<sub>x</sub>) is the character of, A<sub>0</sub>(N<sub>x</sub>) we have that ϕ(x) = ϕ<sub>0</sub>(N<sub>x</sub>), x ∈ G, is the lifted character of A(x).

The lifting operation preserves irreducibility.

### Proof

Suppose that A<sub>0</sub>(N<sub>x</sub>), is a representation of G/N having degree m and that

$$A(x) = A_0(N_x), x \in G,$$

A(x) A(y) = A<sub>0</sub>(N<sub>x</sub>) A<sub>0</sub>(N<sub>y</sub>) = A<sub>0</sub>(N<sub>x</sub> N<sub>y</sub>) = A(N<sub>xy</sub>) = A(xy), which implies that A(x) is a Representation.

From A(x) = A<sub>0</sub>(N<sub>x</sub>) it follows trivially that

$$\phi(x) = \phi_0(N_x),$$

Since A(x) consisting of the same matrices of, A<sub>0</sub>(N<sub>x</sub>), it is clear that A(x) is irreducible if and only if A<sub>0</sub>(N<sub>x</sub>), is irreducible.

An example to illustrate the use of this lifting process in the construction of character table.

**Example 14** (character table for Alternating group of degree four)

	$Z_3$		
$\chi_0$	1	Z	$Z^2$
$\chi_1$	1	1	1
$\chi_2$	1	W	$W^2$
$\chi_3$	1	$W^2$	W

	$S^3$			
	$C_1$	$C_2$	$C_3$	$C_4$
$h^{(4)}$	1	4	4	3
$\chi_1$	1	1	1	1
$\chi_2$	1	W	$W^2$	1
$\chi_3$	1	$W^2$	W	1
$\chi_4$	3	A	B	D

	$A_4$			
	$C_1$	$C_2$	$C_3$	$C_4$
$h^{(4)}$	1	4	4	3
$\chi_1$	1	1	1	1
$\chi_2$	1	W	$W^2$	1
$\chi_3$	1	$W^2$	W	1
$\chi_4$	3	0	0	-1

The alternating group of degree four ( $A_4$ ) is of order 12 and consists of the following elements  
 1, (123), (132), (124), (142), (134), (143), (234), (243), (12)(34), (13)(24), (14)(23).

We first split  $A_4$  into conjugacy classes, namely,

$$\begin{aligned} C_1 &= \{1\} \\ C_2 &= \{(123), (142), (134), (234)\} \\ C_3 &= \{(132), (124), (143), (243)\} \\ C_4 &= \{(12)(34), (13)(24), (14)(23)\} \end{aligned}$$

We know that by Hill (1976)  $A_4$  has four irreducible representations. It so happens that  $U = C_1 \cup C_4$  is a normal subgroup of  $A_4$ . The quotient group  $A_4/U$  is of order three and therefore isomorphic to  $Z_3$  whose character Table 5 is

$$\text{Where } W = e^{2\pi i/3}$$

According to Theorem 13 we can list the three characters of  $Z_3$  all of which are linear to obtain three characters  $\chi_1, \chi_2$  and  $\chi_3$  of  $A_4$ . The kernel being  $U$ . It remains to find  $\chi_4$ .

Since,  $(\chi_1)^2 + (\chi_2)^2 + (\chi_3)^2 = 12$  that is  $\chi_1 = \chi_2 = \chi_3 = 1$  which implies that  $\chi_4 = 3$ .

Let the values of  $\chi_4$  for the classes,  $C_1, C_2$  and  $C_3$  be a, b, c respectively. Thus the character Table 6 for  $A_4$  is,

The values of a, b and c are derived from orthogonality of the columns (character relations of the second kind). Thus it is found that  $a = b = 0$  because  $1 + W + W^2 = 0$  and  $3d + 3 = 0$ . Hence,  $d = -1$  and the complete character Table 7 for is given.

## REFERENCES

- Curtis, C.W. and I. Reiner, 1962. Representation theory of finite Groups and Associative Algebras. John Wiley and Sons' Inc. New York.
- Dornhoff, L., 1971. Group representation theory, Dekker, New York.
- Feit, W., 1971. Characters of finite Groups, New York.
- Hill, V.E., 1976. Group representation and characters. Hafens Press, Macmillian Publishing Co. Inc, New York.
- Isaacs, I.M., 1956. Character theory of finite Group Academic Press New York.
- Ledermann, W., 1976. Introduction to Group characters Cambridge University Press.