

Economic Value of Dried Tomato Pomace in Broiler Chickens Diets

E. Rahmatnejad, M. Bojarpour, M. Islami, J. Fayazi and M. Mamouei
Department of Animal Science,
Ramin University of Agricultural and Natural Resources, Ahvaz, Iran

Abstract: This experiment was conducted to study economic value of Dried Tomato Pomace (DTP) in broiler chickens diets in the form of completely randomized design. For this purpose, 160 days old Ross 308 broilers were distributed into 16 floor pens and reared for 42 days. A basal diet was formulated according to NRC recommendations for starter (0-21 days) and grower (22-42 days) periods. Then, the basal diet was also, supplemented with 8, 16 and 24% DTP, resulting 4 dietary treatments were prepared including control group. The results showed that feed conversion ratio between treatments containing 8 and 16% of DTP compared to control group had no significant difference ($p>0.05$), but the highest (worst) feed conversion ratio were recorded for birds fed diets supplemented with 24% DTP ($p<0.05$). Comparison of the cost for 1 kg meat showed that the use of DTP up to 16% in diet reduce production cost.

Key words: DTP, economic value, broiler, diets, protein

INTRODUCTION

Despite all attempts to set the world population, problem in the human population explosion in developing countries is quite evident. This huge population needs to food and protein after oxygen. Broiler production in Iran developed very fast in the last two decades to become the most important sector in animal production. The major problems that face poultry production particularly broiler industry in Iran are the availability and the cost of feed ingredients. Exploitation of agricultural by-products may make a substantial contribution towards better and more economic feeding of poultry. In view of the shortage and the high costs of protein feeding-stuffs, some of these by-products such as DTP could provide part of the protein needed by poultry. In this respect, it is clear that the country's resources in such products need to be fully developed and thoroughly evaluated nutritionally. DTP one of wet tomato pomace is a mixture of tomato skin, pulp and crushed seeds that remain after the processing of tomato for juice, paste and/or ketchup. The proximate analysis shows that DTP may contain acceptable nutrients such as crude protein, crude fibre, diethyl ether extract, nitrogen free extract (Persia *et al.*, 2003; King and Zeidler, 2004). The seed protein is rich in lysine and can supplement products that are deficient in this amino acid like cereals (Tsatsaronis and Boskou, 1975; Carlson *et al.*, 1981; Liadakis *et al.*, 1995). Be noted that the tomato pomace metabolizable energy content was 2130 kcal kg⁻¹ (NRC, 1988). Manifold studies available about effect of DTP on performance and carcass characteristic broiler

chickens (Dotas *et al.*, 1999; King and Zeidler, 2004; Al-Betawi, 2005; Jafari *et al.*, 2006), but there are limit information about the economic value of DTP in broiler chickens diets. Therefore, the purpose of this study was to investigate the economic value of using DTP in broiler diets.

MATERIALS AND METHODS

The experiment was carried out at the poultry research station, Ramin Research Institute, Faculty of Animal Science and Food industry, Ramin Agricultural and Natural Resources University. The experimental procedure was approved by the Animal Research Committee of the Ramin Agricultural and Natural resources University.

Chemical analysis: In this study, samples of DTP for chemical analysis transmitted to animal science group laboratory and analysis was performed according to AOAC (1984) methods (Table 1) and compared to other finding (Squires *et al.*, 1992; Hassan and Sh, 2004; Hamza, 2001; NRC, 1994, 1988).

Birds, diets and statistical analysis: In this experiment, 160 broiler chickens of the commercial Ross 308 strain were used in a randomized completely design with 4 treatment, 4 replicates in each treatment and 10 birds/replicates and reared on the floor pens for 42 days. A basal diet was formulated as control according to NRC (1994), recommendations for starter (0-21 days) and

Table 1: Chemical composition of tomato pomace (%)

References	Nutrients							
	DM	CP	CF	Ash	EE	NFE	Ca	P
This experiment	90.6	22.0	27.0	0.43	11.9	22.6	0.43	0.63
Squires <i>et al.</i> (1992)	93.7	18.8	31.0	3.60	12.2	28.0	0.41	0.54
Hassan and Sh (2004)	72.9	22.1	29.3	11.10	7.9	31.5	-	-
Hamza (2001)	92.2	22.7	32.4	5.60	12.3	26.3	-	-
NRC (1994)	92.0	23.0	26.0	6.00	-	-	0.43	0.59
NRC (1988)	92.0	23.5	26.4	7.50	10.3	-	0.43	0.60

Table 2: Ingredient composition and calculated analysis of diets

Ingredients	Starter diets DTP (%)				Grower diets DTP (%)			
	0	8	16	24	0	8	16	24
Corn	57.2	56.7	50	44.1	62.6	63.6	58	49
Soybean meal	29.9	27.8	25	24	24.2	19.5	14	15
Tomato pomace	0	8	16	24	0	8	16	24
Wheat bran	5	0	0	0	5	0	0	0
Fish meal	3	3	3	3	5.1	6.1	6.8	5.1
Soybean oil	1.2	1	2.2	3.5	1.7	1	2	3.8
Oyster shell meal	1	1.1	1.1	1.1	1.3	1.3	1.3	1.4
DCP	1.3	1.1	0.9	0.73	0.86	0.39	0	0
Vit and min premix	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5
DL-methionine	0.16	0.17	0.20	0.26	0.09	0.07	0.07	0.15
L-lysine	0.26	0.16	0.14	0.16	0.22	0.08	0.03	0.11
NaHCO ₃	0.14	0.17	0	0	0.11	0	0	0
Salt	0.13	0.18	0.20	0.20	0.14	0.18	0.20	0.19
Coccidiostat	0	0	0	0	0.05	0.05	0.05	0.05
Total	100	100	100	100	100	100	100	100
Nutrient content								
ME (Kcal kg ⁻¹)	2850	2850	2850	2850	2950	2950	2950	2950
Crude protein (%)	20.48	20.48	20.48	20.48	18.44			
Crude fiber (%)	3.88	5.38	7.21	9.04	3.6	4.9	6.7	8.6
Cost (Rail)	4980	4840	4810	4830	4790	4550	4490	4620

Vitamin and mineral provided per kg of diet: Vitamin A, 360000 IU; Vitamin D₃, 800000 IU; Vitamin E, 7200 IU; Vitamin K₃, 800 mg; Vitamin B₁, 720 mg; Vitamin B₆, 400 mg; Vitamin H₂, 40 mg; Vitamin B₁₂, 2640 mg; Vitamin B₃, 4000 mg; Vitamin B₅, 12000 mg; Vitamin B₉, 1200 mg; Vitamin B₁₂, 6 mg; Choline chloride, 200000 mg; Manganese, 40000 mg; Iron, 20000 mg; Zinc, 40000 mg; copper, 4000 mg; Iodine, 400 mg; Selenium, 80 mg

grower (22-42 days) periods. The required amount of unusual feedstuff under study was added to the basal diet so that in addition to the control treatment, three dietary experimental treatments containing 8, 16 and 24% DTP were prepared (Table 2). During the experiment, water and feed were given to the birds *ad libitum* also, base of economic value was the cost for 1 kg broiler meat at the end of experiment, calculations are given:

$$C_f = \frac{(a \times a') + (b \times b')}{a + b}$$

$$C_M = FCR \times C_f$$

Where:

C_b , C_M = Cost of 1 kg feed in 0-42 days and cost of 1 kg meat, respectively

a , a' = Total feed consumption and cost of 1 kg feed in starter period, respectively

b , b' = Total feed consumption and cost of 1 kg feed in grower period, respectively

FCR = Feed Conversion Ratio

All data were analyzed using the One-Way ANOVA procedure of SAS[®] (1998) for analysis of variance. Significant differences among treatments were identified at 5% level by Duncan (1955) multiple range tests.

RESULTS AND DISCUSSION

One of the major problems facing poultry industry particularly broiler production in Iran as well as in most development countries is the availability and cost of feed ingredients. This study is intended to examine the potential use of some unusual feed resources that obtained from agro-industrial by-products such as DTP, which could be used in poultry feeding.

The effect of experimental treatments in broiler chickens diets on cost production in whole period (0-42 days) is given in Table 3. DTP to use in diets significantly influence the feed conversion ratio of broiler chicken ($p < 0.05$). In the other hand, treatment contain 24% DTP had the worst feed conversion ratio due to high fiber content. Buchanan *et al.* (2007), Al-Betawei (2005),

Table 3: Effect of inclusion DTP in broiler diets on cost kg⁻¹ meat production in whole period (0-42 days)

DTP (%)	FCR (g g ⁻¹)	Feed (rial)	Meat (Rial)	Meat (%)
		-----Cost kg ⁻¹ -----		
0	1.91 ^b	4834.70	9234.27	100.0
8	1.97 ^b	4618.00	9097.60	98.5
16	1.94 ^b	4563.00	8988.02	97.3
24	2.30 ^a	4673.25	10748.47	116.3

^{a,b}Means in each column with different superscripts are significantly different (p<0.05)

Malathi and Devegowda (2001) and Bedford (2000) have observed the negative effects of fiber on performance of broiler chickens. As shown in Table 3, the cost kg feed decreased with increasing DTP in diets in compared with control group. The cost/kg feed of treatment containing 16% DTP was lowest, due to the low price of DTP in diet containing 24% had the highest cost in comparison to another diets containing DTP, this diet was low metabolizable energy that compensated by soybean oil (with high cost), therefore, this diet was the most expensive among the other diets containing DTP. In case of cost feed and feed conversion ratio, diets containing 8 and 16% DTP were the best for cost per kg meat production. Finally, using DTP up to 16% in broiler chicks diets had the positive effect on economic value of production.

CONCLUSION

Based on the results of this experiment, at inclusion DTP up to 16% in broiler chickens diets reduce production cost.

ACKNOWLEDGEMENTS

The researchers wish to thank the Dr. K.H. Mirzadeh and Dr. M. Chajji for their cooperation.

REFERENCES

- Al-Betawi, N.A., 2005. Preliminary study on tomato pomace as unusual feedstuff broiler diets. Pak. J. Nutr., 4: 57-63. <http://scialert.net/qredirect.php?doi=pjn.2005.57.63&linkid=pdf>.
- AOAC, 1984. Official Methods of Analysis. 14th Edn. Assoc. Off. Anal. Chem., Arlington, VA. ISBN: 0-935 584-24-2.
- Bedford, M.R., 2000. Exogenous enzymes in monogastric nutrition their current value and future benefits. Anim. Feed Sci. Technol., 86: 1-13. <http://www.oie.int/eng/publicate/RT/2401/24-1%20pdfs/12-madan127-140.pdf>.
- Buchanan, N.P., L.B. Kimbler, A.S. Parsons, G.E. Seidel, W.B. Bryan, E.E. Felton and J.S. Moritz, 2007. The effect of nonstarch polysaccharide enzyme addition and dietary energy restriction on performance and carcass quality of organic broiler chickens. Applied Poult. Res., 16: 1-12. <http://japr.fass.org/cgi/content/short/16/1/1>.
- Carlson, B.L., D. Knorr and T.R. Watkins, 1981. Influence of tomato seed addition on the quality of wheat flour breads. J. Food Sci., 46: 1029-1031, 1042. DOI: 10.1111/j.1365-2621.1981.tb02985.x.
- Dotas, D., S. Zamanidis and J. Balios, 1999. Effect of dried tomato pulp on the performance and egg traits of laying hens. Br. Poult. Sci., 40: 695-697. DOI: 10.1080/00071669987106.
- Duncan, D.B., 1955. Multiple range and F-tests Biometrics, 11: 1-42.
- Hamza, R.G., 2001. Effect of gamma irradiation and enzyme supplementation on the nutritional and biological values of tomato and pea wastes. Ph.D. Thesis, Faculty of Agriculture, Cairo University.
- Hassan, M. and M. Sh, 2004. Nutritional studies on carp fish M.Sc Thesis, Faculty of Agriculture, Moshtohor, Zagazig University, Banha Branch.
- Jafari, M., R. Pirmohammadi and V. Bampidis, 2006. The use of dried tomato pulp in diets of laying hens. Int. J. Poult. Sci., 5 (7): 618-622. <http://scialert.net/qredirect.php?doi=ijps.2006.618.622&linkid=pdf>.
- King, A. and G. Zeidler, 2004. Tomato pomace may be a good source of vitamin E in broiler diets. California Agric., 58 (1): 59-62. <http://calag.ucop.edu/0401JFM/pdfs/pomace.pdf>.
- Liadakis, G.N., C. Tzia, V. Oreopoulou and C.D. Thomopoulos, 1995. Protein Isolation from tomato seed meal, extraction optimization. J. Food Sci., 60: 477-82. DOI: 10.1111/j.1750-3841.1998.00450.pp.x.
- Malathi, V. and G. Devegowda, 2001. *In vitro* evaluation of nonstarch polysaccharide digestibility of feed ingredients by enzymes. National Academy Press, Washington DC. Poult. Sci., 80: 302-305. <http://ps.fass.org/cgi/reprint/80/3/302?maxtoshow=&HITS=10&hits=10&RESULTFORMAT=&fulltext=Malathi+and+Devegowda&andorexactfulltext=and&searchid=1&FIRSTINDEX=0&sortspec=relevance&resource=HWCIT>.
- NRC, 1994. National Research Council. Nutrient Requirements of Poultry. 9th Edn. National Academy Press, Washington, DC. <http://books.nap.edu/openbook.php?isbn=0309048923>.

- NRC (National Research Council), 1988. Nutrient Requirements of Domestic Animals. 1st Edn. National Academy Press, Washington, DC. <http://books.nap.edu/booksearch.php?term=Nutrient+requirements+of+domestic+animals%2C+1988&isbn=0309048923>.
- Persia, M.E., C.M. Parsons, M. Schang and J. Azconat, 2003. Nutritional evaluation of dried tomato seeds. *Poult. Sci.*, 82: 141-146. <http://ps.fass.org/cgi/reprint/82/1/141.pdf>.
- SAS Institute Inc., 1998. SAS/STAT User's Guide. Edn. SAS Institute Inc., Cary, North Carolina.
- Squires, M.W., E.C. Naber and V.D. Toelle, 1992. The effect of heat, water, acid and alkali treatment of tomato cannery wastes on growth, metabolism energy value and nitrogen utilization of broiler chicks. *Poult. Sci.*, 71: 522-529.
- Tsatsaronis, G.C. and D.G. Boskou, 1975. Amino acid and mineral salt content of tomato seed and skin waste. *J. Sci. Food Agric.*, 26: 421-423. DOI: 10.1002/jsfa.2740260406.