

Antimicrobial Properties of the Ethanolic Extracts of *Zingiber officinale* (Ginger) on *Escherichia coli* and *Pseudomonas aeruginosa*

¹K.I. Auta, ¹A.A. Galadima, ¹J.U. Bassey, ¹O.D. Olowoniyi, ²O.O. Moses and ³A.B. Yako

¹Department of Science Laboratory Technology, ²Department of Mathematics and Statistics,
Federal Polytechnic Institute, P.M.B. 01, Nasarawa, Nigeria

³Department of Biological Sciences, Faculty of Natural and Applied Sciences,
Nasarwa State University, Keffi, Nigeria

Abstract: The antimicrobial properties of various extract of *Zingiber officinale* (ginger) against *Escherichia coli* and *Pseudomonas aeruginosa* that are common causes of gastrointestinal track infection were investigated using the Agar and tube diffusion method. The results obtained revealed that etharolic extract of ginger gave the widest zone of inhibition against one out of the two test organisms at the concentration of 20 mg mL⁻¹. However, *Pseudomonas aeruginosa* was more sensitive to the extract. It was also observed that the solvent of extraction and its varying concentrations affected the sensitivity of the two test organisms to the plant extract showing that ginger has antimicrobial activities on the test organisms due to its inhibitory effect, thus confirming its use in folk medicine.

Key words: Antimicrobial properties, *Zingiber officinale*, *Escherichia coli*, *Pseudomonas aeruginosa*, folk medicine, Nigeria

INTRODUCTION

The use of medicinal plants to treat ailment associated with pains is well known through history (Ernst and Pittler, 2000). Such plants can play important role in drug discovery and this study is logical research strategies in the search for new drugs.

Ginger, the underground rhizome of the plant *Zingiber officinale* has been used as a medicinal plant in Asia, India, Jamaica and Nigeria. In China, ginger has been used to aid digestion, treat stomach upset, diarrhea and nausea for >2000 years (Muaz, 1999; Azu and Onyeagba, 2007).

Ginger has a wide range of action on the human body and has been found effective in the treatment of cataract, heart disease, migraines, struck amenorrhea, athletes foot, bursitis, chronic fatigue, cold, flu, coughs, depression, dizziness, fever, arectile difficulties, kidney stones, reynad's disease and viral infection (Peggy, 2006). Ginger has also been historically used to treat inflammation which several scientific studies support through one arthritics that showed ginger to be no better that a placebo or ibuprofen. Research on rats suggests that ginger may be useful for treating diabetes; in the West, powdered derived ginger root is made into capsules and sold in pharmacies for medical use (Ody, 1997; Ohae,

1999). In Venezuela, ginger is pounded into a paste and applied to the abdomen for difficult menstruation. In Costa Rica, it is used in a decoction to relieve throat inflammation and asthma with the addition of honey, it is a valued remedy for coughs and bronchitis and also serves as a sudorific in fever. Its natural diuretic stimulates the kidney to flush out toxins faster. In Panama, it is used to relieve rheumatism.

In Guatemala and Trinidad, it is the best remedy for stomachache, malaria and indigestion the fumes from an infusion in urine are inhaled to relieve head colds. Due to prevalent resistance of micro-organism to drugs and other therapeutic agent, this research is aimed at investigating the potentials of ginger as a medicinal plant that will provide a natural and locally produced antimicrobial agent with an effective rate compared to the standard antibiotics commercially produced against pathogenic micro-organisms.

MATERIALS AND METHODS

Sampling: The test organisms, *Escherichia coli* and *Pseudomonas aeruginosa* were collected from National Institute of Pharmaceutical Research, Abuja. The pure culture was sub-cultured on nutrient agar then preserved in the refrigerator at 4°C until it was required for study.

The plant material *Zingiber officinale* (ginger) was purchased from Nasarawa market and authenticated at the College of Agriculture Lafia, Nasarawa state.

Extraction of plant material (Ginger): The ginger rhizomes were washed with distilled water and allowed to dry (air-day) for 2 days. Extraction was done using the following procedures; crude extraction methods; cold water extraction and ethanolic extraction. About 150 g of fresh ginger was blended into zinc powder and soaked in 100 mL of distilled water for 24 h.

The pulp obtained was left in a clean sterile glass container and shaken vigorously to allow proper extraction and it was filtered using sterile muslin cloth to obtain the filtrate and evaporated to dryness using water bath and stored below ambient temperature.

About 150 g fresh ginger was soaked in 100 mL of 95% ethanol for 24 h to obtain the extract which was evaporated to dryness using water bath and stored as in above 150 g of fresh ginger was blended and the raw juice was extracted after standing in a clean glass container for 24 h, it was extracted using sterile muslin cloth and the extract was evaporated to dryness using water bath stored.

Preparation of McFarland standard: About 0.5 McFarland equivalent turbidity standard was prepared by adding 0.6 mL of 1% barium chloride dehydrate solution ($\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$) to 99.4 mL of 1% sulphuric acid solution (H_2SO_4). A small volume of the turbid solution was transferred to capped tube of the same type that was used to prepare the test and of control inocula. This was stored in the dark at room temperature.

Preparation of inoculum: Inoculums were prepared by direct colony suspension where a small volume of sterile was poured into a test tube to which general colonies of the test organisms taken directly from the plate was emulsified and the suspension adjusted to match the 0.5 McFarland standard which has similar appearance of an overnight broth culture by adding distilled water (Azu and Onyeagba, 2007; Garc and Bruc, 1993).

Antimicrobial screening test: The sensitivity of the test organisms to the extract of *Zingiber officinale* (ginger) was carried out using the Agar and test tube diffusion method. Total 0.02 mL of the suspension was added to an already prepared medium using a glan dropper and a sterile loop was used to spread by streaking the organisms all over the surface of the medium and allow to dry for 5 min. Cups of 6 mm in diameter was made in the agar using sterile cork borer.

Different dilutions of the plant extracts: Different dilutions of the plant extracts in the order of 20, 5 and 1.25 mg mL⁻¹ were prepared, respectively in 5 different test tubes and place in a test tube rack. About 0.3 mg L⁻¹ of erythromycin was also prepared alongside which serves are a positive control and distilled water as a negative control. Each concentration was introduced into each hole on the medium, as well with disc soak with the various concentrations of *Zingiber officinale* extract on the medium and left to stand on the bench for 1 h for proper diffusion. It was incubated at 37°C for 24 h.

RESULTS AND DISCUSSION

The results of the antimicrobial properties of the extracts on the test organisms are shown in Table 1-3. The result of this research indicates that the extracts of *Zingiber officinale* have antimicrobial properties when the extracts were tested on *P. aeruginosa* and *E. coli*. The widest zone of inhibition was obtained with *P. aeruginosa* while *E. coli* showed a little zone of inhibition. The difference in the zone of inhibition may be directly related to the susceptibility of each test organism to the ginger extracts. The factors responsible for this high

Table 1: Sensitivity pattern of *Escherichia coli* and *Pseudomonas aeruginosa* to raw *Zingiber officinale* extract

Concentration (mg mL ⁻¹)	Zone of inhibition diameter (mm)		Positive control (erythromycin)	Negative control (distilled water)
	<i>E. coli</i>	<i>P. aeruginosa</i>		
20	0.0	3.0	24.0	0.0
10	0.0	5.0	24.0	0.0
5.0	0.0	7.0	24.0	0.0
2.5	0.0	0.0	24.0	0.0
1.25	0.0	0.0	0.0	0.0

Table 2: Sensitivity pattern of *Escherichia coli* and *Pseudomonas aeruginosa* to cold water extract *Zingiber officinale*.

Concentration (mg mL ⁻¹)	Zone of inhibition diameter (mm)		Positive control (erythromycin)	Negative control (distilled water)
	<i>E. coli</i>	<i>P. aeruginosa</i>		
20	12	14	26	0.0
10	10	12	24	0.0
5.0	8	9	24	0.0
2.5	5	3	24	0.0
1.25	2	2	20	0.0

Table 3: Sensitivity pattern of *Escherichia coli* and *Pseudomonas aeruginosa* to ethanolic extract *Zingiber officinale*

Concentration (mg mL ⁻¹)	Zone of inhibition diameter (mm)		Positive control (erythromycin)	Negative control (distilled water)
	<i>E. coli</i>	<i>P. aeruginosa</i>		
20	13.0	15	26.0	0.0
10	11.0	11	26.0	0.0
5.0	8.0	9	24.0	0.0
2.5	0.0	5	24.0	0.0
1.25	0.0	3	0.0	0.0

NB: Values are means of duplicate readings

susceptibility of *P. aeruginosa* to the extract may be attributed to the presence of secondary plant metabolites. Also, the positive control in each of the test organism was sensitive given a value of 24 mm for *E. coli* and 20 mm for *P. aeruginosa* in the raw extract of ginger, 24 mm for *E. coli* and 26 mm for *P. aeruginosa* in the ethanolic extract of ginger and 24 mm for *E. coli* and 20 mm for *P. aeruginosa* in the cold water extract of ginger. About 0.5 McFarland given an equivalent approximate density of bacteria 1×10^{-8} cfu 28 (Azu and Onyeagba, 2007).

CONCLUSION

It became clear in this research that the solvent of extraction affected the degree of antimicrobial activity of the extracts. It was observed that the ethanolic extract of *Zingiber officinale* gave the widest zone of inhibition (15 mm) and *P. aeruginosa* gave a zone of inhibition (13 mm) using the concentration of 20 mg mL⁻¹ each. This credit of ethanolic extraction is due to the fact that ethanol is an organic solvent that will dissolve organic compound better (like dissolves like), hence liberates the active component required for antimicrobial activity. It was observed that raw extract had activity only on *P. aeruginosa* and no effect was observed on *E. coli*. The non inhibition of growth in *E. coli* may be due to non liberation of the active constituents of the raw extract. The cold water extract inhibited the growth in both organism

at all concentration. This is due to water ability to liberate the active constituent of the plant. It is note worthy that the antibacterial activities of these plant is depended on the concentration of the extract.

REFERENCES

- Azu, N.C. and R.A. Onyeagba, 2007. Antimicrobial of extract of *Allium Copa* (Onions) and zingiber *Bacillum subtilis*. Int. J. Trop. Med., 3: 1-4.
- Ernst, E. and M.H. Pittler, 2000. Efficacy of ginger for nausea and vomiting: A systematic review of randomized clinical trials. Br. J. Anesthes., 84: 367-371.
- Garc, L.S. and D.A. Bruc, 1993. Diagnostic Medical Phraseology. 2nd Edn., American Society of Microbiology, Washington DC.
- Muaz, I., 1999. Personal communication Ethnobotanist. National Institute of Pharmaceutical Research and Development (NIPRD) 1999, Abuja.
- Ody, P., 1997. The Complete Medicinal Herbs. Dorling Kindersley Ltd., London.
- Ohae, A., 1999. Personal communication taxonomist. National Institute of Pharmaceutical Research and Development (NIPRD) Abuja.
- Peggy, T.F., 2006. Ginger selection and storage information. Marco Polo Revived Interest on Ginger, pp: 1.