



## OPEN ACCESS

### Key Words

Fetal malnutrition, clinical assessment of nutritional

### Corresponding Author

Amit Kumar Singh,  
Department of Pediatrics, All India  
Institute of Medical Sciences Bhopal,  
Madhya Pradesh, India

**Received:** 12 June 2023

**Accepted:** 21 June 2023

**Published:** 3 July 2023

**Citation:** Ankit Yadav, Brijesh Kumar Patel, Mohd Asad Khan and Amit Kumar Singh, 2023. Assessment of Fetal Malnutrition by Various Anthropometric Parameters. Res. J. Med. Sci., 17: 362-375, doi: 10.59218/ makrjms.2023.362.375

**Copy Right:** MAK HILL Publications

## Assessment of Fetal Malnutrition by Various Anthropometric Parameters

<sup>1</sup>Ankit Yadav, <sup>2</sup>Brijesh Kumar Patel, <sup>3</sup>Mohd Asad Khan and <sup>4</sup>Amit Kumar Singh

<sup>1</sup>Department of Pediatrics, PCMS and RC Bhopal, Ankit, India

<sup>2</sup>Department of Pediatrics, Birsa Munda Government Medical College Shabdol, Madhya Pradesh, India

<sup>3</sup>Department of Pediatrics, LN Medical College and JK Hospital, Bhopal, India

<sup>4</sup>Department of Pediatrics, All India Institute of Medical Sciences Bhopal, Madhya Pradesh, India

### ABSTRACT

Fetal malnutrition (FM) is a clinical state, characterized by intrauterine loss of or failure to acquire normal amount of fat and muscle mass. Assessment of fetal malnutrition should be included in the evaluation of all newborns regardless of the classification of their weight for Gestational Age (GA), as birth weight alone is a poor indicator of nutritional status at birth. Nutritional status at birth can be assessed by using various anthropometric parameters [weight, length, head circumference (HC), chest circumference (CC), midarm circumference (MAC), proportionality indices [Body mass index (BMI), Ponderal index (PI), Kanawati Index (MAC/HC)] and Clinical Assessment of Nutritional status score (CAN score). Present study was a cross sectional, observational and analytical study, conducted in post natal ward and NICU of department of Paediatrics of Peoples College of Medical Sciences and Research Centre, Bhopal during one and half year (1st December, 2018 to 31st may, 2020) of study period. Mean value of all anthropometric parameters i.e. birth weight, length, head circumference, chest circumference and mid arm circumference was significantly lower in those with FM as compared to those without FM ( $p < 0.001$ ).

## INTRODUCTION

Fetal malnutrition (FM) is a clinical state, characterized by intrauterine loss of or failure to acquire normal amount of fat and muscle mass. This term was coined by Scott and Usher in 1963<sup>[1]</sup>.

Assessment of fetal malnutrition should be included in the evaluation of all newborns regardless of the classification of their weight for gestational age (GA), as birth weight alone is a poor indicator of nutritional status at birth<sup>[2,3]</sup>.

Various terminologies used for describing intra uterine malnutrition includes small for gestational age (SGA), intra uterine growth restriction (IUGR) and placental insufficiency. Although these terms are used synonymously with FM, but both are quite different as they do not assess the accumulation of subcutaneous fat and muscle mass in the fetal body<sup>[2]</sup>. Also, they do not take account of genetic and ethnic variations amongst different populations<sup>[4]</sup>.

It is important to recognize FM early in neonates as there is a high incidence of neonatal morbidity and mortality as well as long term neurological sequelae associated with it. Studies have shown that children with FM are more likely to have lower IQ scores, require higher need of special education, have neurologic disabilities, intellectual disabilities, learning disorders or seizures in late childhood as compared to children without FM<sup>[5,6]</sup>. Neurologic alterations may be aggravated by events like hypoglycemia or feeding difficulties happening during the neonatal period that are more common in FM babies. Furthermore, FM is associated more frequently with cardiovascular, endocrine and metabolic disorders during late childhood and adolescent age group<sup>[6,7]</sup>.

The incidence of low birth weight (LBW) babies (<2500 g) continues to be high in India at about 18% in contrast to 5-7% in developed countries<sup>[8,9]</sup>. High incidence of LBW babies in India is due to the neglected health and education of females, teenage marriages and pregnancies, frequent pregnancies, maternal malnutrition, bad obstetric history, pregnancy induce hypertension (PIH), anemia and infections<sup>[10]</sup>.

Nutritional status at birth can be assessed by using various anthropometric parameters [weight, length, head circumference (HC), chest circumference (CC), midarm circumference (MAC), proportionality indices [Body mass index (BMI), Ponderal index (PI), Kanawati Index (MAC/HC)] and Clinical Assessment of Nutritional status score (CAN score)<sup>[3,11,12]</sup>.

Importance of addressing this hidden FM is emphasized because of its potentially serious effects on multiple organ systems. And, if it can be appropriately diagnosed and treated in babies at risk, it decreases morbidity, improve survival and lessens long term sequelae.

Assessment in perinatology (Maternal-Fetal medicine) uses anthropometry as an essential tool to monitor growth and evaluate the nutritional and functional (circulatory and respiratory) status of newborns (NB). It is an economical, non-invasive and easy-to- execute tool that can improve understanding of growth patterns and their variations<sup>[13]</sup>.

Maternal Urinary tract infection and genital tract infection during pregnancy were associated with LBW baby or preterm delivery, urinary tract infection also associated with pregnancy induce hypertension (PIH), anemia and amnionitis which also leads to LBW or preterm LBW newborn<sup>[14]</sup>.

Various instruments have been used to identify children with malnutrition. In older children and adults, Body Mass Index (BMI) is used as the gold standard in determining body proportions and adiposity for the screening of malnutrition. In neonates also as in older children's various criteria have been used to identify and classify FM. The most common criterion used is birth weight. Researchers have argued that birth weight alone may not reflect the true state of nutrition in utero<sup>[11,15,16]</sup>.

Recording of birth weight in villages has continued to be major problem in our country, therefore search was continued for an alternative measurement which could replace birth weight recording to screen for high-risk infants.

Body mass index (BMI) values for gestational age in all percentiles shows a steady increase up to 38 weeks, level off up to 40 weeks followed by slight decrease at 42 weeks in both gender, results show direct correlation between gestational age and BMI and can provide useful reference to assess intra uterine proportional growth<sup>[11]</sup>.

## MATERIALS AND METHODS

**Aim:** Present study was a cross sectional, observational and analytical study, conducted in post-natal ward and NICU of department of Pediatrics of Peoples College of Medical Sciences and Research Centre, Bhopal during one and half year (1st December, 2018 to 31st may, 2020) of study period.

### Objectives:

- To assess fetal nutritional status by assessing newborn using selected anthropometric indices.
- To compare the assessment of fetal nutritional status using CAN score with selected anthropometric indices.

**Source of data:** New born babies were recruited from People's Hospital, which is a allied hospital of Peoples College of Medical Sciences and Research Center

Bhopal, a tertiary referral center, getting patients from all socio-economic groups. A total of four hundred eleven new born babies were included in this study.

**Inclusion criteria:**

- All full-term (37 completed weeks of gestation) newborns as assessed by Modified Ballard score system

**Exclusion criteria:**

- New-borns <37 completed weeks of gestation
- New-borns with congenital anomalies
- Babies born to mothers with Gestational Diabetes Mellitus

**Sample size:** All full term (37-42 weeks of gestational age) newborns delivered at People’s Hospital, Bhopal during the study period were included in the study.

**Methodology:** Written consent (Annexure-III) was taken from parents before enrolling the babies in the study. Babies who fulfilled the exclusion criteria were excluded from the study. All babies were examined thoroughly as per pre-defined proforma (Annexure-II). GESTATIONAL AGE (GA) was assessed by using Modified Ballard scoring system within first 24-48 hrs of life; babies who were found to be full term i.e., 37-42 weeks by Ballard scoring system were enrolled in the study:

- Equipment/Instruments required
  - Electronic weighing machine (Fig. 1):
  - Electronic weighing machine
  - Infantometer (Fig. 2)
  - Flexible non stretchable tape (Fig. 3)

**Variables studied in the study:**

- Mode of delivery
- Singleton/Multiple gestation
- Gestational Age as assessed by Modified Ballard Score
- Birth weight
- Length
- Head circumference
- Chest Circumference
- Mid Arm circumference
- Mid Arm circumference/ Head circumference
- Ponderel Index (PI) 11. Body mass index
- CAN score
- High risk factors in mothers

**Birth weight:** Nude birth weight at birth was measured to the nearest 10 g using electronic weighing scale IN Fig. 4. Birth weight was plotted on Lubchencho chart for weight for gestational age Fig. 5 and baby was classified as SGA (wt/gestational Age less than 10th percentile), AGA (between 10th to 90th percentile) and LGA (more than 90th percentile) (Fig. 4 and 5).

**Length:** Crown to Heel Length was measured to the nearest 0.1cm using an infantometer and length was plotted on length for gestational age Lubchencho chart (Fig. 6 and 7).

**Head circumference:** Occipito-frontal circumference was taken as the largest circumference of the skull using a flexible non stretchable tape to the nearest of 0.1 cm (Fig. 8). Head circumference was also plotted on Lubchencho chart and classified as head circumference less than 10th centile and more than 10th centile (Fig. 9).

**Chest circumference:** Was taken at the level of nipple using flexible non-stretchable tape to the nearest 0.1 cm (Fig. 10).

**Mid arm circumference (MAC):** Measured in the left arm, at a point midway between tip of the acromion and the olecranon process using a flexible non stretchable tape to the nearest 0.1 cm. MAC less than -2SD was taken as abnormal (Table 1).



Fig. 1: Electronic weighing machine



Fig. 2: Infantometer



Fig. 3: Non stretchable measuring tape



Fig. 4: Measurement of weight




Fig. 5: Lubchenko chart: weight for gestational age



Fig. 6: Measurement of length by infantometer

- Mid arm circumference/head circumference Ratio (MAC/HC) (Kanawati Index): A cut off value of  $<0.25$  was used in this study to define malnutrition (Fig. 11)<sup>[4]</sup>.




Fig. 7: Lubchencho chart: length for gestational age



Fig. 8: Measurement of head circumference




Fig. 9: Lubchencho chart: head circumference for gestational age



Fig. 10: Measurement of chest circumference



Fig. 11: Measurement of mid arm circumference

- Ponderal index (PI):

$$PI = \text{Weight (g)} \times 100 / \text{Length (cm)}^{-3}$$

- Ponderal index less than  $2.2 \text{ g cm}^{-3}$  was considered as an indicator of malnutrition<sup>[17]</sup>.
- Body mass index (BMI):

$$BMI = \text{Weight (Kg)} / \text{Length (m)}^2$$

BMI was plotted on BMI charts for different gestational age available for female and male child separately BMI less than 10th centile was considered abnormal (Fig. 12 and 13)<sup>[11]</sup>.

**Statistical analysis:** All data compiled in Microsoft Excel and the data analysis was performed using softwares IBM SPSS ver. 20 and MED CALC 19.5. Frequency distribution and cross tabulation were used to prepare the tables. Quantitative data is expressed as mean and standard deviation whereas categorical data is expressed as number and percentage. Means were

Table 1: Circumference measurement

| Gestational age(week) | N  | MAC (cm)  | MAC/HC    | Birth weight (g) |
|-----------------------|----|-----------|-----------|------------------|
| 25-26                 | 5  | 4.90±0.7  | 0.22±0.02 | 838±249          |
| 27                    | 7  | 5.25±0.3  | 0.22±0.01 | 1022±143         |
| 28                    | 10 | 5.50±0.5  | 0.23±0.02 | 1064±193         |
| 29                    | 6  | 5.70±0.4  | 0.23±0.02 | 1159±132         |
| 30                    | 8  | 6.00±0.7  | 0.23±0.02 | 1307±159         |
| 31                    | 6  | 6.40±1.0  | 0.23±0.03 | 1399±308         |
| 32                    | 14 | 7.00±0.5  | 0.24±0.02 | 1658±231         |
| 33                    | 12 | 7.00±0.8  | 0.24±0.02 | 1750±281         |
| 34                    | 6  | 8.30±0.5  | 0.27±0.01 | 2291±348         |
| 35                    | 12 | 8.10±0.6  | 0.26±0.01 | 2299±308         |
| 36                    | 12 | 8.30±0.6  | 0.26±0.02 | 2364±329         |
| 37                    | 6  | 9.50±0.7  | 0.28±0.02 | 2901±194         |
| 38                    | 22 | 9.50±0.7  | 0.28±0.01 | 3054±348         |
| 39                    | 22 | 9.70±0.9  | 0.28±0.02 | 3076±398         |
| 40                    | 36 | 10.10±0.6 | 0.29±0.02 | 3261±311         |
| 41                    | 11 | 10.20±0.6 | 0.29±0.02 | 3447±337         |
| 42                    | 5  | 10.60±0.5 | 0.30±0.01 | 3383±186         |

Mid-arm circumference measurement, MAC to HC ratio and Birth weight of infant of 25-42 weeks of gestational age expressed in Mean±SD

compared using One Way Anova test. Percentage and numbers were compared using Chi square test and level of significance was considered at 5% Gestational age (GA) assessment by new Ballard scoring system (Fig. 14).




Fig. 12: BMI for gestational age male newborn




Fig. 13: BMI for gestational age in female new born

Fetal malnutrition (FM) is a clinical state defined by intra uterine loss or failure to acquire normal amount of subcutaneous fat and muscle mass and is independent of birth weight and gestational age. During the study period of one and half year from 1st December 2018 to 31st may 2020 we have enrolled 411 full terms new born babies and nutrition status of each new born is assessed at 24-48 hrs of life by using CAN Score and other anthropometric parameters.

Table 2 shows distribution according to gender. Mean (SD) of weight, length, head circumference, chest circumference and mid arm circumference (MAC) and proportionality indices like BMI (weight in kg/length in m<sup>2</sup>, PI (weight in grams/length in cm<sup>3</sup>×100) and Kanawati Index (MAC/HC) are calculated (Table 2 and 3).

Mean weight, length, head circumference, chest circumference, MAC, MAC/HC, PI, BMI and CAN Score

Fig. 14: new ballard score for gestational age assessment

Table 2: Distribution of babies according to gender

| Gender      | Frequency | Percentage |
|-------------|-----------|------------|
| Male        | 217       | 52.80      |
| Female      | 194       | 47.20      |
| Grand total | 411       | 100.00     |

Table 3: Anthropometric indicators of growth in study cohort (n = 411)

| Variable                  | Mean    | SD      |
|---------------------------|---------|---------|
| Weight (g)                | 2774.25 | 405.780 |
| Length (cm)               | 47.54   | 1.640   |
| Head circumference (cm)   | 33.25   | 1.120   |
| Chest circumference (cm)  | 30.95   | 1.360   |
| MAC (cm)                  | 9.51    | 0.700   |
| Kanawati Index (MAC/HC)   | 0.28    | 0.016   |
| PI (g cm <sup>-3</sup> )  | 2.56    | 0.290   |
| BMI (kg m <sup>-2</sup> ) | 12.20   | 1.400   |
| CAN score                 | 26.94   | 2.31    |

SD: Standard deviation

is 2774.25 g, 47.54, 33.25, 30.95, 9.51, 0.28, 2.56 g cm<sup>-3</sup>, 12.20 kg m<sup>-2</sup> and 26.94, respectively (Table 4).

The distribution of babies into well-nourished and malnourished groups using cut-offs of various parameters. It shows that most commonly used anthropometric parameter, weight for gestational age identified 36 (8.8%) babies as malnourished, while BMI cutoff (<10th centile) could identify 210 (51.09%) babies as malnourished. CAN score (<25) used as gold

Table 4: Distribution of babies into well-nourished and malnourished groups based on different parameters

| Parameters            | No of patients | Percentage |
|-----------------------|----------------|------------|
| <b>CAN score</b>      |                |            |
| <25                   | 76             | 18.5       |
| ≥25                   | 335            | 81.5       |
| Total                 | 411            | 100        |
| <b>MAC/HC</b>         |                |            |
| <0.25                 | 23             | 5.6        |
| ≥ 0.25                | 388            | 94.4       |
| Total                 | 411            | 100        |
| <b>Ponderal index</b> |                |            |
| <2.2                  | 56             | 13.6       |
| ≥2.2                  | 355            | 86.4       |
| Total                 | 411            | 100        |
| <b>Weight for GA</b>  |                |            |
| <10th centile         | 36             | 8.8        |
| ≥10th centile         | 375            | 91.2       |
| Total                 | 411            | 100        |
| <b>BMI</b>            |                |            |
| <10th centile         | 210            | 51.09      |
| ≥10th centile         | 201            | 48.91      |
| <b>Length for GA</b>  |                |            |
| <10th centile         | 16             | 3.90       |
| ≥10th centile         | 395            | 96.1       |
| Total                 | 411            | 100        |

standard in present study could identify 76 (18.5%) babies as malnourished (Fig. 15 and Table 5).

Table 6 shows the association between anthropometric parameters and CAN Score cutoff of <25. We found strong association between all the






Fig. 15: Distribution of babies into well nourished and malnourished groups using cut-offs of various parameters




Fig. 16: AUC for the birth weight for determining FM (0.891 with p-value of <0.001)




Fig. 17: AUC for MAC for determining FM (0.855 with p-value of <0.001)

variables with CAN Score cutoff of <25. Out of 411 babies, CAN score cutoff of <25 identified 76 (18.5%) as malnourished and 335 (81.5%) as well nourished. Weight for gestational age identified 36 (8.8%) babies as SGA and 375 (91.2%) as AGA; 46 (12.3%) babies who were identified AGA by weight for gestational age were classified as malnourished by CAN score. MAC/HC ratio cutoff of <0.25 identified 23 (5.6%) babies as malnourished and 388 (94.4%) babies as well nourished; 55 (14.2%) babies who were identified as well nourished by MAC/HC ratio were classified as malnourished by CAN score.

In present study nutritional status of new born babies were studied with various anthropometric parameters, proportionality indices and CAN score. Comparison of various parameters in identifying FM with CAN score (CAN score <25 in identifying FM taken as gold standard) was done. It was found that sensitivity was highest with BMI (89.5%) followed by MAC for age (46.4%), weight for GA (39.5%) and PI (38.2%). Specificity for FM was found to be highest with MAC/HC (99.6%) followed by length for GA (98.8%), HC for GA (98.8%), weight for GA (98.2%) and PI (91.9%). Specificity of BMI was found to be lower (57.6%) as compared to other parameters (Table 7).

In present study nutritional status of new born is assessed by CAN score and compared with selected anthropometric parameters. Comparison of Mean±SD of anthropometric data of babies done with and without FM (as by CAN score cutoff of <25 as gold standard). It was found that birth weight, length, HC, CC and MAC all were significantly lower in those with FM as compared to those without FM (p<0.001). (Table 8).

Comparison of anthropometric parameters with and without FM (FM is defined here as CAN score <27). When modified CAN score Cutoff was considered, 132 (32.5%) babies were found to be malnourished as compared to 76 (18.5%) babies with cutoff of <25 (Fig. 16-21). Analysis shows a strong strength of association (p-value<0.001) between modified CAN

Table 5: Comparison of anthropometric parameters with and without FM with CAN score cutoff of <25

|                       | CAN score  |                  |              | p-value |
|-----------------------|------------|------------------|--------------|---------|
|                       | FM (<25)   | Without FM (≥25) | Total        |         |
| <b>Weight for GA</b>  |            |                  |              |         |
| <10 centile           | 30 (83.4%) | 6 (16.6%)        | 36 ( 8.8% )  | <0.001  |
| ≥10th centile         | 46 (12.3%) | 329 (87.7%)      | 375 (91.2%)  |         |
| Total                 | 76 (18.5%) | 335 ( 81.5%)     | 411 ( 100%)  |         |
| <b>MAC/HC</b>         |            |                  |              |         |
| <0.25                 | 21 (91.4%) | 2 (8.6%)         | 23 (5.6%)    | <0.001  |
| ≥0.25                 | 55 (14.2%) | 333 (85.8%)      | 388 (94.4%)  |         |
| Total                 | 76 (18.5%) | 335 (81.5%)      | 411 (100%)   |         |
| <b>Ponderal index</b> |            |                  |              |         |
| <2.2                  | 29 (51.7%) | 27 (48.3%)       | 56 (13.6%)   | <0.001  |
| ≥2.2                  | 47 (13.2%) | 308 (86.8%)      | 355 (86.4%)  |         |
| Total                 | 76 (18.5%) | 335 (81.5%)      | 411 (100%)   |         |
| <b>BMI</b>            |            |                  |              |         |
| <10th centile         | 68 (32.4%) | 142 (67.4%)      | 210 (51.09%) | <0.001  |
| ≥10th centile         | 8 (3.9%)   | 193 (96.1%)      | 201 (48.91%) |         |
| Total                 | 76 (18.5%) | 335 (81.5%)      | 411 (100%)   |         |
| <b>Length for GA</b>  |            |                  |              |         |
| <10th centile         | 12 (75%)   | 4 (25%)          | 16 (3.9%)    | <0.001  |
| ≥10th centile         | 64 (16.2%) | 331 (83.8%)      | 395 (96.1%)  |         |
| Total                 | 76 (18.5%) | 335 (81.5%)      | 411 (100%)   |         |




Fig. 18: AUC curve for BMI for determining FM (0.837 with p-value of <0.001)




Fig. 20: AUC cure for length for GA in determining FM (0.749 with p-value of <0.001)




Fig. 19: AUC curve for PI in determining FM (0.761 with p-value of <0.001)




Fig. 21: AUC curve for MAC/HC for determining FM (0.714 with p-value of <0.001)

Table 6: Comparison of various parameters in identifying FM with CAN score cutoff of <25

| Parameters  | Weight for GA | PI    | MAC/HC | BMI  | Length for GA | HC for GA | MAC for GA |
|-------------|---------------|-------|--------|------|---------------|-----------|------------|
| Sensitivity | 39.50         | 38.20 | 16.20  | 89.5 | 15.80         | 11.80     | 46.40      |
| Specificity | 98.20         | 91.90 | 99.60  | 57.6 | 98.80         | 98.80     | 97.70      |
| PPV         | 83.30         | 51.80 | 88.30  | 32.5 | 75.00         | 69.20     | 87.40      |
| NPV         | 87.70         | 86.80 | 83.40  | 96.0 | 83.80         | 83.20     | 92.30      |
| LR          | 81.95         | 38.42 | 19.24  | 61.8 | 25.57         | 16.71     | 18.55      |

Table 7: Comparison of Mean±SD of anthropometric data of babies with and without FM

| Parameters               | CAN score      |                  | p-value |
|--------------------------|----------------|------------------|---------|
|                          | With FM (<25)  | Without FM (>25) |         |
| Birth weight (g)         | 2325.53±274.40 | 2870.90±358.46   | <0.001  |
| Length (cm)              | 46.29±1.85     | 47.85±1.46       | <0.001  |
| Head circumference (cm)  | 32.04±1.04     | 33.50±0.98       | <0.001  |
| Chest circumference (cm) | 29.76±1.47     | 31.24±1.18       | <0.001  |
| MAC (cm)                 | 8.89±0.56      | 9.67±0.57        | <0.001  |

Table 8: Comparing anthropometric parameters with and without FM (FM is defined by modified CAN score cutoff of <27)

|                       | CAN score   |                  |             | p-value |
|-----------------------|-------------|------------------|-------------|---------|
|                       | FM (<27)    | Without FM (>27) | Total       |         |
| <b>Weight for GA</b>  |             |                  |             |         |
| <10 cent              | 32 (88.9%)  | 4 (11.1%)        | 36 (8.8%)   | <0.001  |
| >10th cent            | 100 (26.7%) | 275 (73.3%)      | 375 (91.2%) |         |
| Total                 | 132 (32.1%) | 279 (67.9%)      | 411 (100%)  |         |
| <b>MAC/HC</b>         |             |                  |             |         |
| <0.25                 | 22 (95.7%)  | 1 (4.3%)         | 23 (5.6%)   | <0.001  |
| ≥0.25                 | 110 (28.4%) | 278 (71.6%)      | 388 (94.4%) |         |
| Total                 | 132 (32.1%) | 279 (67.9%)      | 411 (100%)  |         |
| <b>Ponderal index</b> |             |                  |             |         |
| <2.2                  | 38 (67.9%)  | 18 (32.1%)       | 56 (13.6%)  | <0.001  |
| ≥2.2                  | 94 (26.5%)  | 261 (73.5%)      | 355 (86.4%) |         |
| Total                 | 132 (32.1%) | 279 (67.9%)      | 411 (100%)  |         |
| <b>BMI</b>            |             |                  |             |         |
| <10th cent            | 112 (53.3%) | 98 (46.7%)       | 210 (51.1%) | <0.001  |
| ≥10th cent            | 20 (9.95%)  | 181 (90.05%)     | 201 (48.9%) |         |
| Total                 | 132 (32.1%) | 279 (67.9%)      | 411 (100%)  |         |

Table 9: AUC curve for various parameters with 95% confidence interval

| Variables    | AUC   | SE    | 95% confidence interval (CI) |
|--------------|-------|-------|------------------------------|
| CAN score    | 1.000 | 0.000 | 1.000-1.000                  |
| Birth weight | 0.891 | 0.020 | 0.853-0.930                  |
| MAC          | 0.855 | 0.026 | 0.805-0.905                  |
| BMI          | 0.837 | 0.024 | 0.785-0.880                  |
| PI           | 0.761 | 0.031 | 0.701-0.822                  |
| Length       | 0.749 | 0.033 | 0.684-0.814                  |
| MAC/HC       | 0.714 | 0.033 | 0.702-0.831                  |

SE: Standard error

score cutoff of <27 and various anthropometric variables (Table 9) (weight for gestational age, PI, BMI and MAC/HC).

## DISCUSSIONS

In accordance to our study that shows better specificity and PPV of MAC/HC over PI, Georgieff *et al.*<sup>[12]</sup> also studied the accuracy of PI and MAC/HC ratio for detecting babies who are likely to be symptomatic because of aberrant intrauterine growth and found that MAC/HC ratio is more accurate than PI for evaluation of potentially symptomatic newborn who suffered abnormal fetal growth. Similar to present study, study done by Soundarya *et al.*<sup>[18]</sup>, identified newborn as either well-nourished or malnourished by PI, MAC/HC, BMI and CAN score. Few babies who were found to be normal by PI, MAC/HC ratio and BMI (25 out of 222 in PI, 42 out of 219 in MAC/HC and 11 out of 179 in BMI group) were found to be malnourished by CAN score. The study also concluded that FM is best identified by CAN Score, BMI is best screening tool for

FM and when coupled with PI will identify most normally nourished newborns<sup>[19]</sup>. Present study observed 18.5% babies to be malnourished by using CAN score and 13.6% by PI which is less than observed by Abhay kumar Dhanorkar (32.29% and 24.48% respectively)<sup>[20]</sup>.

To classify nutritional status of new born, PI has been used by various investigators<sup>[21,22]</sup>. PI relies on the principle that length is spared at the expense of weight during period of acute conditions; whereas, weight and length velocities are proportionately impaired in chronic insults. Therefore, using PI alone as a method of nutritional assessment can misclassify the babies. The other drawback of PI is that any error in calculating length is cubed in the calculation of PI<sup>[23]</sup>.

Many cities now have multi-ethnic population and application of weight standard alone may be inappropriate in studying nutritional status. To overcome this problem, many investigators<sup>[24,25]</sup> studied MAC/HC ratio in identifying nutritional status and concluded that this ratio shows no intra ethnic variation and can be used as screening test for identifying babies whose growth is retarded, even when their weight does not fall below 10th centile. However, in chronic in-utero insult, head circumference is also reduced because of proportionate growth retardation; therefore such babies are missed by MAC/HC ratio. Similarly, babies with hydrocephalus may give falsely low reading even

when they are normally nourished. CAN score can identify FM in these babies too who are missed by PI and MAC/HC ratio.

In a study by Ezenwa *et al.*<sup>[26]</sup>, the nutritional status of full term new born baby was done using CAN score and compared it with PI, BMI and MAC/HC. FM was identified in 14.5, 10.3, 13.1 and 2.8% of babies using CAN score, PI, BMI and MAC/HC ratio respectively. Out of FM babies identified by CAN score, PI, MAC/HC and BMI identified FM in 19.5, 12.3 and 53.7% babies respectively which shows that BMI was most sensitive anthropometric index for detecting FM. In present study CAN score identified FM in 18.5% babies whereas MAC/HC, PI, weight for gestational age, BMI and length for gestational age identified FM in 5.6, 13.6, 8.8, 51.09 and 4.4% newborn babies respectively. Out of FM identified by CAN score, MAC/HC, PI, weight for gestational age, length for gestational age and BMI identified FM in 16.2, 38.2, 39.2, 15.8 and 89.5% new born babies, respectively. Thus, present study also shows that BMI has maximum sensitivity in identifying FM.

In present study we assessed the FM by preparing Receiver Operating Characteristic curve (ROC curve) and Area Under Curve (AUC). The ROC-AUC analysis showed that AUC for birth weight, MAC, BMI, PI, length and MAC/HC for determining FM was 0.891, 0.855, 0.837, 0.761, 0.749 and 0.714 respectively with a p-value of <0.001. AUC was highest for birth weight followed by MAC and BMI. This highlight that all these three parameters have higher sensitivity in determining FM (p-value <0.001). Similarly a study done by Sen J *et al* observed that AUC for birth weight (0.796, 95% CI 0.741-0.850) followed by MAC (AUC 0.776, 95% CI 0.721-0.831) can be considered to be the best surrogate anthropometric measures of low birth weight<sup>[27]</sup>. However, they have not used the CAN score as a measure of FM in their study. This highlights that birth weight and BMI are strongly associated with FM, however if coupled with CAN score can provide better assessment for FM.

In a study done by Mohan *et al.*<sup>[23]</sup>, assessment of nutritional status of new born was done using CAN score cutoff of <25 as well as <21 along with other anthropometric parameters. In this study more number of babies were clustered between 21 and 24, so they have taken modified CAN score cutoff as <21<sup>[28]</sup>. On the contrary, in present study large number of babies were clustered between 27 to 29 CAN score, so we selected <27 as modified cutoff score. The explanation behind considering this modified cutoff is that in a community large proportion of babies cannot be called as abnormal. In present study, on using modified CAN score cutoff <27, the percentage of FM went up from 18.5-32.11%. This signifies that when we are using modified CAN score cutoff of <27, we are able to detect more malnourished newborn than using cutoff of <25<sup>[29-36]</sup>.

#### Limitations of the study:

- Being a hospital based and single centered study, it is difficult to extrapolate the results found in large community
- In present study, observations were done by single observer only therefore, inter-observer variations could not be calculated.

#### Strength of the study:

- Sufficiently large sample size that includes neonates belonging to all socio- economic strata was the main strength of this study

**Implication of study:** Malnourished neonates missed by most commonly used parameter i.e. Weight for GA (> 10th centile) can be detected by CAN score as malnourished. Proper nutritional counselling of mother and long term follow up of these babies can prevent neuro- developmental adverse outcomes of these babies.

Fetal malnutrition (FM) is a clinical state characterized by intrauterine loss of or failure to acquire normal amount of fat and muscle mass. The term was coined by Scott and Usher in 1963.

It is important to recognize FM in babies because of high incidence of neonatal morbidity and mortality and long term neurological sequelae associated with it. Existing terminologies for describing intra uterine malnutrition includes, small for gestational age (SGA), intra uterine growth restriction (IUGR) and placental insufficiency. None of these terminologies are synonymous with FM as they do not assess the accumulated subcutaneous fat and muscle mass in the fetal body. Also, they do not take account of genetic and ethnic variations amongst different populations.

#### RESULTS

Present study is designed to identify the nutritional status by clinical assessment of newborn using CAN score and compare it with selected other anthropometric indices. It was a cross sectional, observational and analytical study and included all full term new born babies (gestational age assessed by Modified Ballard score system) delivered at People's Hospital from 1st December 2018 to 31st may, 2020, which is a allied hospital of Peoples College of Medical Sciences and Research Centre Bhopal, a tertiary referral centre getting patients from all socio-economic groups. New born less than 37 completed weeks gestation, congenital anomalies and babies born to mother with gestational Diabetes mellitus (Infant of diabetic mother) were excluded from study.

Data was collected within 24-48 hrs of life of newborn (n = 411) after obtaining written consent from parents. Birth weight was recorded using digital electronic weighing machine, length was measured

using infantometer, Head circumference, chest circumference and mid arm circumference was measured using non stretchable measuring tape. Proportionality indices like Kanawati Index (MAC/HC), PI and BMI was calculated for each new born baby and clinical assessment of new born using CAN score (based on superficial physical findings) was done for each newborn. Data was compiled using Microsoft Excel and analyzed using softwares SPSS® Version 20 and MED CALC 19.5.

The results can be summarized under following points:

- The mean value of weight, length, head circumference, chest circumference, MAC, MAC/HC, PI, BMI and CAN Score in subjects included in present study was 2774.25 g, 47.54, 33.25, 30.95, 9.51, 0.28, 2.56 g cm<sup>-3</sup>, 12.20 kg m<sup>-2</sup> and 26.94, respectively
- Male to female ratio was 1.12: 1
- Fetal malnutrition (FM) was found in 18.5% babies as assessed by the CAN score of <25
- Out of 411 babies, 76 (18.5%) were with FM (CAN score <25), 23 (5.6%) had MAC/HC ratio <0.25, 56 (13.6%) had Ponderal Index of <2.2, weight for GA was
- <10th centile in 36 (8.8%) and BMI was <10th centile in 51.09% babies.
- Weight for GA (p<0.001), MAC/HC (p<0.001), PI (p<0.001), BMI (p<0.001) and length for GA (P<0.001) all were found to have significant association in predicting the fetal malnutrition when CAN score of <25 was taken as the cut off value
- Mean value of all anthropometric parameters i.e. birth weight, length, head circumference, chest circumference and mid arm circumference was significantly lower in those with FM as compared to those without FM (p<0.001)
- Maternal risk factors such as PIH, anaemia, maternal infection and poor socioeconomic status have significant association with FM, whereas birth spacing (year), BMI of mother and hypothyroidism were not significantly associated with the presence of FM
- With modified CAN score cutoff of <27, percentage of FM among AGA babies goes up from 12.26% to 26.66%.
- The area under the curve (AUC) for the birth weight (0.891), MAC (0.855), BMI (0.837), PI (0.761), length (0.749) and MUAC/HC (0.714) with p-value of <0.001 was observed. AUC was highest for birth weight followed by MAC and BMI. This highlight that all these three parameters have good sensitivity and specificity for identifying FM (p<0.001)

## CONCLUSIONS

Fetal malnutrition (FM) is a major problem in developing countries like India as compared to developed countries. FM is a clinical state defined by intrauterine loss or failure to acquire normal amount of subcutaneous fat and muscle mass and is independent of birth weight and gestational age. Previous studies have shown that commonly used method of classifying babies on basis of weight for GA as SGA, AGA and LGA may miss to diagnose FM in some of the cases who are affected late in third trimester. Various methods are used for assessment of nutritional status of new born but each method has its own limitations.

## REFERENCES

1. Scott, K.E. and R. Usher, 1966. Fetal malnutrition: Its incidence, causes, and effects. *Am. J. Obstet. Gynecol.*, 94: 951-963.
2. Metcalf, J., 1994. Clinical assessment of nutritional status at birth: Fetal malnutrition and SGA are not synonymous. *Pediatr. Clin. North. Am.*, 41: 875-891.
3. Sankhyan, N., V.K. Sharma and S. Singh, 2009. Detection of fetal malnutrition using "CAN score". *Indian J. Pediatr.*, 76: 903-906.
4. Meadows, N.J., J. Till, A. Leaf, E. Hughes, B. Jani and V. Larcher, 1986. Screening for intrauterine growth retardation using ratio of mid-arm circumference to occipitofrontal circumference. *Br. Med. J.*, 292: 1039-1040.
5. Hill, R.M., W.M. Verniaud, R.L. Deter, L.M. Tennyson and G.M. Rettig *et al.*, 1984. The effect of intrauterine malnutrition on the term infant.. *Acta Paediatrica*, 73: 482-487.
6. Adebami, O.J. and J.A. Owa, 2008. Comparison between CANSORE and other anthropometric indicators in fetal malnutrition. *Indian J. Pediatr.*, 75: 439-442.
7. Fok, T.F., K.L. Hon, P.C. Ng, E. Wong and H.K. So *et al.*, 2008. Use of anthropometric indices to reveal nutritional status: Normative data from 10, 226 Chinese neonates. *Neonatology*, 95: 23-32.
8. IIPS, 2017. National family health survey (NFHS-4). International Institute for Population Sciences Deonar, <http://rchiips.org/NFHS/NFHS-4Reports/India.pdf>
9. WHO., 2004. Low Birthweight: Country, Regional and Global Estimates. World Health Organization., ISBN-10: 9280638327, Pages: 31.
10. Deodhar, J. and R. Jarad, 1999. Study of the prevalence of and high risk factors for fetal malnutrition in term newborns. *Ann. Trop. Paediatrics*, 19: 273-277.
11. Brock, R.S., M.C. Falcão, C. Leone 2008. Body mass index values for newborns according to gestational age. *Nutr Hosp* 23: 487-492.

12. Georgieff, M.K., S.R. Sasanow, U.M. Chockalingam and G.R. Pereira, 1988. A comparison of the mid-arm circumference/head circumference ratio and ponderal index for the evaluation of newborn infants after abnormal intrauterine growth. *Acta Paediatrica*, 77: 214-219.
13. Altman, D.G. and F.E. Hytten, 1989. Intrauterine growth retardation: Let's be clear about it. *BJOG: An Int. J. Obstet. Gynaecol.*, 96: 1127-1128.
14. Lakhute, S.V., V.V. Kendre, J.V. Dixit, A. Nagaonkar and L.T. Chinte, 2016. Comparative study of CANSCORE with anthropometry in the assessment of fetal malnutrition at tertiary care hospital. *Int. J. Recent Trends Sci. Technol.*, 18: 316-319.
15. Amarendra, M. and M. Yoganand, 2017. Comparison of clinical assessment of nutritional status (CAN) score with other methods in the assessment of fetal malnutrition. *Int. J. Contemp. Pediatr.*, 4: 713-718.
16. Vikram, S., A. Prashant and K. Nutan, 2012. Detection of fetal malnutrition by CAN score at birth and its comparison with other methods of determining intrauterine growth. *Indian J. Clin. Prac.*, 22: 576-582.
17. Clifford, S.H., 1954. Postmaturity with placental dysfunction. *J. Pediatr.*, 44: 1-13.
18. Soundarya, M., A. Basavaprabhu, K. Raghuvveera, B. Baliga and B. Shivanagaraja *et al.*, 2012. Comparative assessment of fetal malnutrition by anthropometry and can score. *Iran.J. Pediatr.*, 22: 70-76.
19. Bhargava, S.K., 1991. Perspectives in child health in India. *Indian Pediatr*, 28: 1403-1410.
20. Shastry, C.K.R. and R.B. Poornima, 2015. Anthropometric measurements of newborns. *Int. J. Contemp. Pediatr.*, 2: 85-89.
21. Villar, J. and J. Belizán, 1982. The relative contribution of prematurity and fetal growth retardation to low birth weight in developing and developed societies. *Am. J. Obstet. Gynecol.*, 143: 793-798.
22. Bhargava, S.K., S. Ramji, A. Kumar, M. Mohan, J. Marwah and H.P. Sachdev, 1985. Mid-arm and chest circumferences at birth as predictors of low birth weight and neonatal mortality in the community. *BMJ*, 291: 1617-1619.
23. Mohan, M., S.R. Prasad, H.K. Chellani and V. Kapani, 1990. Intrauterine growth curves in north Indian babies: weight, length, head circumference and ponderal index. *Indian Pediatr.*, 27: 43-51.
24. Schieve, L.A., A. Handler, R. Hershov, V. Persky and F. Davis, 1994. Urinary tract infection during pregnancy: Its association with maternal morbidity and perinatal outcome. *Am. J. Public Health*, 84: 405-410.
25. Ezenwa, B., E. Iroha, V. Ezeaka and M.C. Egri-Okwaji, 2016. Comparative study of clinical assessment of nutritional status score and proportionality indices in the assessment of fetal malnutrition in term newborns. *Niger. Med. J.*, 57: 124-128.
26. Sasanow, S.R., M.K. Georgieff and G.R. Pereira, 1986. Mid-arm circumference and mid-arm/head circumference ratios: Standard curves for anthropometric assessment of neonatal nutritional status. *J. Pediatr.*, 109: 311-315.
27. Mehta, S., A. Tandon, T. Dua, S. Kumari and S.K. Singh *et al.*, 1998. Clinical assessment of nutritional status at birth. *India. Pediatr.*, 35: 423-428.
28. Rao, M.R., N. Balakrishna and K.V. Rao, 1999. Suitability of CANSCORE for the assessment of the nutritional status of newborns. *Indian J. Pediatr.*, 66: 483-492.
29. Kashyap, L. and R. Dwivedi, 2006. Detection of fetal malnutrition by clinical assessment of nutritional status score (can score) at birth and its comparison with other methods of determining intrauterine growth. *Pediatr. Oncall J.*, 3: 9-12.
30. Sajjadian, N., H. Shajari, F. Rahimi, R. Jahadi and M.G. Barakat, 2011. Anthropometric measurements at birth as predictor of low birth weight. *Health*, 3: 752-756.
31. Waghmare, P., D.N. Balpande and B.B. Lakhkar, 2012. Assessment of fetal malnutrition by CAN score. *Pediatr. Oncall. J.*, 9: 6-8.
32. Tiwari, A.K., D. Bandyopadhyay and B. Saha, 2018. Anthropometric indices and clinical assessment of nutrition score (CAN Score) in detection of fetal malnutrition at birth. *Int. J. Pediatr. Res.*, 5: 175-179.
33. Ballard, J.L., J.C. Khoury, K. Wedig, L. Wang, B.L. Eilers-Walsman and R. Lipp, 1991. New Ballard score, expanded to include extremely premature infants. *The J. Pediatr.*, 119: 417-423.
34. Lubchenco, L.O., C. Hansman, M. Dressler and E. Boyd, 1963. Intrauterine growth as estimated from liveborn birthweight data at 24 to 42 weeks of gestation. *Pediatrics*, 32: 793-800.
35. Hendrix, N. and V. Berghella, 2008. Non-placental causes of intrauterine growth restriction. *Seminars Perinatology*, 32: 161-165.
36. Brodsky, D. and H. Christou, 2004. Current concepts in intrauterine growth restriction. *J. Intensive Care Med.*, 19: 307-319.