



OPEN ACCESS

Key Words

Uric acid, sepsis

Corresponding Author

S. Sivakumar,
Department of Emergency Medicine
Sree Mookambika Institute of
Medical Sciences Kanyakumari,
Tamil Nadu, India

Author Designation

¹Associate Professor

²Unior Resident

Received: 20th February 2025

Accepted: 10th March 2025

Published: 21st May 2025

Citation: V. Ravishankar and S. Sivakumar, 2025. Estimation of Uric Acid Level for the Assessment of the Severity of Diseases with Clinically Diagnosed Sepsis. Res. J. Med. Sci., 19: 5-8, doi: 10.36478/makrjms.2025.4.5.8

Copy Right: © 2025. V. Ravishankar and S. Sivakumar. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Estimation of Uric Acid Level for the Assessment of the Severity of Diseases with Clinically Diagnosed Sepsis

¹V. Ravishankar and ²S. Sivakumar

¹Department of Emergency Medicine, Sree Mookambika Institute of Medical Sciences Kanyakumari, Tamil Nadu, India

²Department of Emergency Medicine Sree Mookambika Institute of Medical Sciences College Kanyakumari, Tamil Nadu, India

Abstract

The incidences of sepsis and septic shock depend on how acute organ dysfunction and infection are defined as well as on which data sources are studied. Hence this study was conducted to bring out the correlation between hyperuricemia in clinically diagnosed sepsis patients and morbidity and mortality. Total of 150 patients was included in the study. For the uric acid estimation of the included patients, blood samples were taken from the patient. For the biochemical estimation of the uric acid level the collaboration was done with the department of emergency medicine at sree mookambika college of medical sciences. Among the 150 study participants, 64 patients had elevated uric acid levels which constitutes about 42.7%, whereas 86 patients constituting 57.3% had normal uric acid levels. It can be inferred that among the study population, patients had type 2 diabetes mellitus as the most common comorbidity at 40%. The most prevalent comorbidities among the patients with hyperuricemia were diabetesmellitus type 2 and type 1, decompensated liver disease and cerebrovascular accident. Patients without any comorbidities about 36% of the study population also developed sepsis. This study demonstrates that Serum Uric acid may be potentially used as a marker of severity of illness as well as predictor of mortality and morbidity in patients with clinically diagnosed sepsis in the IMCU. This study recommends further studies on a large basis to confirm the observations.

INTRODUCTION

The incidences of sepsis and septic shock depend on how acute organ dysfunction and infection are defined as well as on which data sources are studied. Disparate estimates come from administrative data, prospective cohorts with manual case identification and large electronic health record databases^[1]. Organ dysfunction is often defined by the provision of supportive therapy in which case epidemiological studies count the "treated," rather than the actual, incidence. In the United States, recent cohort studies using administrative data suggest that upwards of 2 million cases of sepsis occur annually^[2,3]. In the past 20 years, research has revealed that infection can cause multiple organ dysfunction but without a measurable inflammatory excess [i.e., without the systemic inflammatory response syndrome (SIRS)]. In fact, both pro and anti-inflammatory responses are present along with significant changes in other pathways. Septic shock is defined as a complication of sepsis resulting in derangements in circulatory and metabolic pathways in the body. Allantoin is the end product of purine metabolism in animals whereas uric acid in human beings^[4,5]. Purines can be endogenous or exogenous. Purines are nitrogenous compounds found in the body as well as food. Uric acid passes through the liver, enters the blood stream and most of it excreted in urine. Some uric acid is degraded in the body after reaction with oxidants.⁶ Over the last ten years, strong association has been found between atherosclerosis, hypertension, hyperinsulinemia and chronic kidney disease and uric acid^[7]. Oxidative stress is found out by the presence of elevated serum uric acid which is a poor prognostic sign in case of patients with sepsis as multi organ dysfunction occurs as a result of high oxygen free radicals. Increased levels of serum uric acid causes acute activation of many transcription factors in patients with severe infection and is a poor prognostic sign in case of severe infection^[8]. Chronic conditions are also associated with elevated serum uric acid. Hence this study was conducted to bring out the correlation between hyperuricemia in clinically diagnosed sepsis patients and morbidity and mortality.

MATERIALS AND METHODS

The present prospective study was done in the medical college and the associated hospital. The study was done over the period of one year. Total of 150 patients were included in the study. The inclusion and exclusion criteria followed were as follows:

Inclusion Criteria:

- Age >18 years.
- Admission to emergency ICU with a working diagnosis of sepsis.

Exclusion Criteria:

- Patients denying consent.
- Pregnant females.
- Known case of kidney disease.
- Patients who have already been in ICU in an outside facility for more than 24 hrs.
- Patients who are known case of gout.
- Patients on drugs causing hyperuricemia. For the uric acid estimation of the included patients; blood samples were taken from the patient. For the biochemical estimation of the uric acid level the collaboration was done with the department of biochemistry, medical college and hospital. The present study was conducted over a period of one year in the IMCU of Sree mookambikai Medical College and Hospital among 150 patients who were >18 years old and admitted in the ICU with a clinical diagnosis of sepsis based on the quick SOFA (qSOFA) score.

Once the patient met the inclusion criteria, the written consent was signed and obtained from the study participants. Clinical proforma for the study including demography was meticulously collected from the study participants. Basic vitals at admission heart rate, respiratory rate, blood pressure, oxygen saturation were recorded. Thorough general and systemic examination performed. Quick SOFA score is based on three parameters, Glasgow coma scale, systolic blood pressure and respiratory rate, assigning one point each to: 1. Low blood pressure SBP \leq 100 mmHg, 2. High respiratory rate (\geq 22 breaths per minute) and 3. GCS $<$ 15. Blood samples were then obtained for uric acid, urea, creatinine, complete blood count, serum electrolytes and chest x-ray was taken. The patient's creatinine at admission to IMCU was taken as the baseline value. Hyperuricemia was defined as value $>$ 7 mg/dl in males and females. Outcome of the sepsis event was classified as either death or discharge from the intensive care unit. Statistical analysis was done using the statistical package for social sciences (SPSS). Different statistical methods were used as appropriate. Mean \pm SD was determined for quantitative data and frequency for categorical variables. The independent t-test was performed on all continuous variables. The normal distribution data was checked before any t-test. The Chi-Square test was used to analyze group difference for categorical variables. A p-value $<$ 0.05 was considered significant.

RESULTS AND DISCUSSIONS

In the present study population, the median age was found to be 59.13 years. There were 6.67% who were under 30 years, 56% belonged to 30-65 years age group and contributed to the maximum number of individuals, followed by 37.33% in the \geq 65 years age

group. Our study population had as light male preponderance at 52%. Uric Acid: Among the 150 study participants, 64 patients had elevated uric acid levels which constitutes about 42.7%, whereas 86 patients constituting 57.3% had normal uric acid levels. It can be inferred that among the study population, patients had type 2 diabetes mellitus as the most common comorbidity at 40%. The most prevalent comorbidities among the patients with hyperuricemia were diabetes mellitus type 2 and type 1, decompensated liver disease and cerebrovascular accident. Patients without any comorbidities about 36% of the study population also developed sepsis. The study participants were found to develop acute kidney injury, acute respiratory distress syndrome as the major complications. Our study aims at understanding the correlation between hyperuricemia and the secondary end points like AKI and ARDS. One of the end points of this study is the outcome of patients status with regards to sepsis and its relation to hyperuricemia. In this study, it was found that out of the 64 patients with hyperuricemia, 46 had expired which constitutes 71.9% and 18 patients were discharged which is 28.1%. However, this difference was found to be not statistically significant.

Table 1: Distribution of Hyperuricemia Among Various Age Groups

Age group	Frequency	Uric acid > 7 mg/dl	Uric acid < 7 mg/dl
<30	10	4	6
31-65	84	36	48
>65	56	24	32
Total	150	64	86

In this prospective cohort study, we report that elevated uric acid levels on arrival to the ICU in patients with sepsis are associated with a poor prognosis; that is, an increased risk for AKI, ARDS, marks an increased duration of stay in the ICU. Sepsis is a condition of increased pro inflammatory cytokines and oxidative stress thereby increases the antioxidants in the body to counterbalance. This altered level of antioxidant defence leads to immune dysfunction and poor outcomes. In a systemic inflammatory response, both endothelial cells and neutrophils are activated to release oxygen derived free radicals^[9,10]. Uric acid increases in response to acute oxidative stress. Uric acid formation may even provide a significant antioxidant defense mechanism against nitration by peroxynitrite in rat heart during hypoxia. Hence uric acid is believed to be an important marker of oxidative stress. The mechanisms for increased uric acid are not well understood^[11]. Development of AKI during sepsis increases patient morbidity, predicts higher mortality, has a significant effect on multiple organ functions, is associated with an increased length of stay in the intensive care unit and hence consumes considerable healthcare resources. The first important finding of our

study is that hyperuricemia is associated with AKI in patients with early sepsis. When AKI develops, then it causes poor prognosis. For instance, the immediate operative and postoperative death rate after CVS surgery ranges between 1-2%, this rises to 10-38% if AKI develops and to >50% if dialysis is needed^[12]. Usually the patients with sepsis are a very complex subset of population who have MODS and bad prognosis and are usually very sick patients. Hemodynamic changes, changes in the functional capacity of the heart and liver, exposure to multiple medications and numerous other factors cause development of AKI in these patients. Among these factors, uric acid can also contribute to development of AKI. The mechanism by which uric acid causes AKI can range from indirect injury secondary to the release of vasoactive mediators and oxidative stress to crystal induced direct tubular toxicity. Uric acid induced renal vasoconstriction due to catecholamine release, activation of renin-angiotensin system, release of pro-inflammatory markers, oxidative stress and decreased nitric oxide levels, in turn causes AKI. Khosla *et al.* have revealed a decrease in plasma nitrites (metabolites of NO) in rats with hyperuricemia by allopurinol. Zoccali^[12] have demonstrated a correlation between high uric acid levels and dysfunction of endothelium. Various pro-inflammatory markers such as MCP and CRP are increased as a response to inflammation induced by uric acid. Uric acid stimulates an inflammatory response via increasing various pro-inflammatory markers such as MCP and CRP^[13,14]. Hence uric acid may be an early marker of impending AKI in patients with sepsis and could be used to predict the risk for AKI in septic patients. This further raises the question of whether the treatment of hyperuricemia in early sepsis could potentially decrease the risk for AKI. Increased uric acid levels play a role not only in the occurrence of AKI but also in the progression of CKD. Uric acid levels are increased in subjects with renal disease as the result of reduction in GFR and renal urate excretion. In our study we found that although there was a high incidence of ARDS noted in this septic patient population, there was no statistically significant association of hyperuricemia with ARDS. Thus although uric acid levels may be used to predict the severity of illness, duration of stay in ICU and risk for AKI, it was not significant enough to predict the incidence of ARDS. This could potentially be due to the small patient population that we had for our study, especially since increasing uric acid levels have been reported by Nagaya *et al.* to correlate with clinical severity of primary pulmonary hypertension and has an independent association with long-term mortality of patient with primary pulmonary hypertension. This was most likely due to the small sample size.

Regarding the outcome of septic patients, though there was a slight increase in mortality among the hyperuricemic individuals with sepsis than those with normal Uric acid levels, it was statistically significant to prove the point. One statistically significant end point was the correlation between mechanical ventilation and ARDS and AKI. It was statistically significant to see increased mechanical ventilation among patients with ARDS.

CONCLUSION

This study demonstrates that Serum Uric acid may be potentially used as a marker of severity of illness as well as predictor of mortality and morbidity in patients with clinically diagnosed sepsis in the ICU. This study recommends further studies on a large basis to confirm the observations.

REFERENCES

1. Martin C.M., F. Priestap, H. Fisher, R.A. Fowler and D.K. Heyland *et al.*, 2009. A prospective, observational registry of patients with severe sepsis: The Canadian Sepsis Treatment and Response Registry*. *Crit. Care Med.*, Vol. 37: 10.1097/CCM.0b013e31819285f0.
2. Jolley R.J., K.J. Sawka, D.W. Yergens, H. Quan, N. Jetté and C.J. Doig., 2015. Validity of administrative data in recording sepsis: A systematic review. *Crit. Care*, Vol. 19: 10.1186/s13054-015-0847-3.
3. Rhee C., M.S. Jentzsch, S.S. Kadri, C.W. Seymour and D.C. Angus *et al.*, 2019. Variation in Identifying Sepsis and Organ Dysfunction Using Administrative Versus Electronic Clinical Data and Impact on Hospital Outcome Comparisons*. *Crit. Care Med.*, Vol. 47. 10.1097/ccm.0000000000003554.
4. Ronco, C., M. Bonello, V. Bordoni, Z. Ricci, V. D'Intini, R. Bellomo and N.W. Levin, 2004. Extracorporeal Therapies in Non-Renal Disease: Treatment of Sepsis and the Peak Concentration Hypothesis. *Blood Purif.*, Vol. 22: 10.1159/000074937.
5. Rubio P.E.A., R.B. Molina, P.E.A. Ávila, A.G. Mora and C.A.G. López., 2019. Infective Endocarditis: Inflammatory Response, Genetic Susceptibility, Oxidative Stress, and Multiple Organ Failure. *Infective Endocarditis*, Vol. 0. 10.5772/intechopen.84908.
6. Barrett K.E., S. Boitano, S.M. Barman and H.L. Brooks., 2010. Ganong's review of medical physiology twenty.
7. Pasalic D., N. Marinkovic and L. Feher-Turkovic., 2012. Uric acid as one of the important factors in multifactorial disorders-facts and controversies. *Biochemia Med.*, Vol. 22: 10.11613/bm.2012.007.
8. Akbar S.R., D.M. Long, K. Hussain, A. Alhajhusain and U.S. Ahmed *et al.*, 2015. Hyperuricemia: An Early Marker for Severity of Illness in Sepsis. *Int. J. Nephrology*, Vol. 2015. 10.1155/2015/301021.
9. Victor V., M. Rocha, J. Esplugues and M. Fuente., 2005. Role of Free Radicals in Sepsis: Antioxidant Therapy. *Curr. Pharm. Des.*, Vol. 11: 10.2174/1381612054864894.
10. El-Shebiny E., S. Daif, S. Shoeib, Y. Fathi and E. Zahran., 2022. Serum uric acid level as a prognostic factor in sepsis outcome. *The Egypt. J. Chest Dis. Tuberculosis*, Vol. 71. 10.4103/ejcdt.ejcdt_3_20.
11. Lappas M., U. Hiden, G. Desoye, J. Froehlich, S.H.D. Mouzon and A. Jawerbaum., 2011. The Role of Oxidative Stress in the Pathophysiology of Gestational Diabetes Mellitus. *Antioxidants and Redox Signaling*, Vol. 15: 10.1089/ars.2010.3765.
12. Zarjou A. and A. Agarwal., 2011. Sepsis and Acute Kidney Injury. *J. Am. Soc. Nephrology*, Vol. 22: 10.1681/ASN.2010050484.
13. Bhargavi S.N., 2020. A Study on Hyperuricemia as an Early Marker for Severity of Illness in Sepsis in IMCU of a Tertiary Care Centre., 2020. Government Vellore Medical College, Vellore., Vol.
14. Kocyigit I., M.I. Yilmaz, O. Orselik, M.H. Sipahioglu and A. Unal *et al.*, 2013. Serum Uric Acid Levels and Endothelial Dysfunction in Patients with Autosomal Dominant Polycystic Kidney Disease. *Nephron Clin. Pract.*, Vol. 123: 10.1159/000353730.